

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 505 680 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

- (45) Date of publication of patent specification: **26.04.95** (51) Int. Cl.<sup>6</sup>: **C07K 7/02, C07K 14/655, A61K 38/31**
- (21) Application number: **92101196.1**
- (22) Date of filing: **27.01.92**

- (54) **Octapeptide or heptapeptide derivatives, a process for preparing them as well as medicaments containing these compounds and the use of them.**

- (30) Priority: **25.01.91 HU 27291**
- (43) Date of publication of application:  
**30.09.92 Bulletin 92/40**
- (45) Publication of the grant of the patent:  
**26.04.95 Bulletin 95/17**
- (84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU NL PT SE**
- (56) References cited:  
**EP-A- 0 203 031**  
**EP-A- 0 215 171**  
**EP-A- 0 277 419**  
**EP-A- 0 298 732**  
**EP-A- 0 389 180**

Proc. Natl. Acad. Sci. USA, 86(6), pages  
2003-2007 (1986)

Greenstein & Winitz, Chemistry of amino acids,  
Wiley, London, 1961, vol. 1, pages 5-6

- (73) Proprietor: **BIOSIGNAL Kutato-Fejlesztő Kft.**  
**Spahl u. 8**  
**H-1022 Budapest (HU)**
- (72) Inventor: **Kéri, György, Dr.**  
**Modori u. 4**  
**H-1021 Budapest (HU)**  
Inventor: **Mező, Imre, Dr.**  
**Hevesi Gy. u. 40**  
**H-1156 Budapest (HU)**  
Inventor: **Horváth, Aniko, Dr.**  
**Ibolya u. 6**  
**H-1038 Budapest (HU)**  
Inventor: **Vadász, Zsolt, Dr.**  
**Gödöllői u. 167/a**  
**H-1141 Budapest (HU)**  
Inventor: **Teplán, István, Dr.**  
**Mihályffy E. u. 35**  
**H-1026 Budapest (HU)**  
Inventor: **Balogh, Agnes**  
**Titel u. 8-10**  
**H-1141 Budapest (HU)**  
Inventor: **Csuka, Orsolya, Dr.**  
**Mártonhegyi u. 22/b**  
**H-1127 Budapest (HU)**

**EP 0 505 680 B1**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

**Current opinion in therapeutic patents,**  
2(12), Dec. 1992, pages 1993-1994

**Biochemical and biophysical research com-**  
**munications, 191(2), 1993, pages 681-687**

Inventor: **Bökönyi, Gyöngyi**  
**Kazai u. 23**

**H-1031 Budapest (HU)**

Inventor: **Szőke, Balázs, Dr.**  
**Lőrinc pap tér 3**

**H-1088 Budapest (HU)**

Inventor: **Horváth, Judit, Dr.**  
**Asztalos J. u. 2/19**

**H-7624 Pécs (HU)**

Inventor: **Idel, Miklos, Dr.**  
**Kárpát u. 16**

**H-1133 Budapest (HU)**

Inventor: **Seprődi, János, Dr.**  
**Szilágyi u. 30**

**H-1028 Budapest (HU)**

⑦ Representative: **Beszédes, Stephan G., Dr.**  
**Patentanwalt**  
**Postfach 11 68**  
**D-85201 Dachau (DE)**

## Description


The invention is concerned with novel octapeptide or heptapeptide amide derivatives, a process for preparing them as well as medicaments containing these compounds and the use of them.

5 Somatostatin, a cyclic tetradecapeptide (SRIF), being an inhibitor of secretion of the growth hormone (GH) was originally isolated from the hypothalamus [Brazeau et al.: Science 179, 77 (1973)]. Somatostatin has a very broad spectrum of biological effects, participates in a high number of biological processes and in the majority of cases it plays a role of an inhibitory factor (it inhibits e.g. the release of prolactin, insulin, glucagon, gastrin, secretin and cholecystoquinine) [S. Reichlin: Somatostatin, N. Eng. J. Med. 309, 1495 and 1536 (1983)].

10 One of the most important effects of somatostatin being a growth-inhibiting factor consists in its capability to influence various forms of pathological cell growth. It is well known from the literature that it exerts an inhibitory action on the growth of cancerous cells [A. V. Schally: Cancer Rs. 48, 6977 (1988); Taylor et al.: Biochem. Biophys. Res. Commun. 153, 81 (1988)]. Likely, somatostatin exerts its antagonizing action on growth factors related to cancerous processes. It has been shown by recent investigations that somatostatin and some somatostatin analogues are capable to activate the tyrosine phosphatase enzyme which antagonizes the effect of tyrosine kinases playing a very important role in the tumorous transformation [A. V. Schally: Cancer Res. 48, 6977 (1988)]. The importance of tyrosine kinases is supported by the fact that the majority of oncogens codes for tyrosine kinase and the major part of the growth factor 15 receptors is tyrosine kinase [Yarden et al.: Ann. Rev. Biochem. 57, 443 (1989)].

Native somatostatin has a very short duration of effect *in vivo* since it is rapidly inactivated by endo- and exopeptidases. A high number of novel analogues have been prepared in order to enhance the duration of effect, biological activity and selectivity of this hormone. Most of the active analogues contain a cycle and a peptide chain which is shorter than the original one. The first cyclic hexapeptide showing the whole 20 effects of somatostatin was synthesized by Veber et al. [Nature 292, 55 (1981)]. As a continuation newer and more effective cyclic hexa- and octapeptides have been synthesized which possess the whole spectrum of effects of somatostatin [Veber et al.: Life Sci. 34, 1371 (1984); Murphy et al.: Biochem. Biophys. Res. Commun. 132, 922 (1985); Cai et al.: Proc. Natl. Acad. Sci. USA 83, 1896 (1986)].

Moreover from EP-A 203 031 there have been known biologically active lysine containing octapeptides 25 of the formula



35

wherein

A represents an L, D or DL amino-acid selected from the group consisting of Ala, Val, Phe, p-Cl-Phe, Trp, Pro, Ser, Thr, Glu, Gly, Beta Ala, Abu, N-Me Ala, 5-F-Trp, 5-Br-Trp, 5-Cl-Trp, their acetylated derivatives or a pharmaceutically acceptable acid addition salt thereof;

40

B represents an L, D or DL amino acid amide selected from the group consisting of Thr NH<sub>2</sub>, Val NH<sub>2</sub>, Pro NH<sub>2</sub>, HO-Pro NH<sub>2</sub>, Ser NH<sub>2</sub>, Tyr NH<sub>2</sub>, Trp NH<sub>2</sub>, 5-F-Trp NH<sub>2</sub>, For-Trp NH<sub>2</sub>, Ala NH<sub>2</sub>, Gly NH<sub>2</sub>, Me Ala NH<sub>2</sub>;

X represents L-Phe or L-Tyr,

45

Y represents L-Thr or L-Val;

Z is L, D or DL-5-F-Trp, 5-Br-Trp, 5-Cl-Trp, 5-I-Trp or D-Trp; and


C'' and C' represent L or D Cys, Abu, Asp or Lys; provided that where C' is Cys, C'' is also Cys and where C' or C'' are other than Cys, C'' is different from C' and is other than Cys and the pharmaceutically acceptable acid addition salts thereof; are useful as agents for inhibiting 50 the release of growth hormone, for the treatment of gastrointestinal disorders and for therapy of certain cancers and the management of diabetes.

The problem underlying to the invention is to create novel somatostatin analogues showing a more advantageous and/or more selective pharmacological action, particularly in treating mammals, including 55 man, suffering from tumour disease, in comparison to that of the known compounds, a process for preparing them as well as medicaments containing these compounds and the use of them.


Surprisingly this has been solved by the invention.

The invention is based on the recognition that the effect of octapeptide and heptapeptide analogues of somatostatin can be strengthened when the amino acid in 1-position is replaced by a substituent of strongly hydrophobic character and having a structure inhibiting the activity of xopeptidases. This substitution can preferably be combined with the replacement of the amino acid in 3-position by various aromatic or heterocyclic amino acids; or with the replacement of the amino acid in 4-position by an aromatic D-amino acid, aromatic aminocarboxylic acid, aminosuccinimide or a  $\beta$ -aspartyl group, optionally substituted on its  $\alpha$ -carboxyl group; or with the replacement of the amino acid in 6-position by an amino acid, bearing an alkyl or aralkyl side chain, or substituted derivatives thereof or proline or  $\beta$ -alanine; or with omission of the amino acid in 6-position; or with replacement of the amino acid in 8-position by an aromatic amino acid or a ring-substituted derivative thereof or by threonine.


A further basis of the invention is the recognition that, when an anthranilyl substitution, for example, in compound of formula




m-aminobenzoyl substitution, for example, in compound of formula



35 or aminosuccinyl substitution, for example, in compound of formula



is used, the compound has not to be cyclized since these replacements inhibit in some cases the free steric rotation of a given peptide (US patent 4,758,552; HU patent 194,280); otherwise the rotation is prevented by the cyclized form, too, for example, in compound of formula



Thus a subject matter of the invention are octapeptide or heptapeptide amide derivatives of the general formula

5

10

15

20

**25**

30

35

40


45

50

55 wherein

X<sub>1</sub> stands for an aromatic D-amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a D-phenylglycyl, p-hydroxyphenylglycyl, o-aminobenzoyl, m-aminobenzoyl, D-tetrahydroisoquinolylcarbonyl or sarcosylalanyl; or a

5




10

in which latter

- $\text{A}_1$  means the rest of an aromatic, heterocyclic or aliphatic amine having a primary or secondary amino group or means a primary or secondary aromatic, heterocyclic or aliphatic amino group or means an alkoxy, aryloxy or aralkyloxy group;  
 $\text{X}_2$  represents an aromatic amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a histidyl group;  
 $\text{X}_3$  means a D-tryptophyl, o-aminobenzoyl, m-aminobenzoyl, aspartyl, D-aspartyl,  $\beta$ -aspartyl,  $\beta$ -D-aspartyl or aminosuccinyl group; or a

20




25

30

in which latter

- $\text{A}_1$  is as defined above;  
 $\text{X}_4$  stands for an amino acid rest bearing an alkyl or aralkyl side chain, or a derivative thereof substituted by a hydroxyl or methyl group in  $\beta$ -position; or a prolyl or  $\beta$ -alanyl group; or a valence bond;  
 $\text{X}_5$  means a heterocyclic amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a threonyl group;  
 $\text{R}_1$  represents hydrogen or a

40



45

in which latter

- $\text{A}_2$  means hydrogen or a  $\text{C}_{1-4}$  alkyl group, optionally substituted by halogen;  
 $\text{R}_2$  means hydrogen, -S- or a -S-acetamidomethyl, phenyl or p-hydroxyphenyl group;  
 $\text{R}_3$  stands for hydrogen or hydroxyl;  
 $\text{R}_4$  means a phenyl or p-hydroxyphenyl group; or -S- or a -S-acetamidomethyl group when the meaning of  $\text{R}_2$  is the same group; or a methyl group when  $\text{R}_3$  is hydroxyl, in case of  $\text{R}_2$  and  $\text{R}_4$  each being -S- one bond each of them are united to a sole bond representing a bridge between  $\text{R}_2$  and  $\text{R}_4$   
 and  
 $n$  is 1, 2, 3 or 4,  
 with the provisos that  
 a) if  
 $\text{X}_1$  stands for D- $\beta$ -Nal or D-Phe, optionally substituted by halogen or a hydroxy group,  
 $\text{X}_2$  means Tyr or Phe

50

55

- $X_3$  represents D-Trp,  
 $X_5$  stands for Trp or Thr and  
 $R_2$  and  $R_4$  each are -S-, one bond each of them being united to a sole bond representing a bridge  
 between  $R_2$  and  $R_4$   
 5  $X_4$  is different from Val, Thr or Ser,  
 b) if  
 $X_4$  stands for Val or Thr or Ser and  
 $X_1$  means an aromatic D-amino acid residue or a derivative thereof ring-substituted by halogen or  
 hydroxy  
 10  $X_2$  represents an aromatic amino acid rest or a derivative ring-substituted by halogen or hydroxy  
 at least one of the following conditions is fulfilled  
 $\alpha$ )  
 $X_3$  is different from D-Trp  
 or  
 15  $\beta$ )  
 $R_2$  and  $R_4$  together have a meaning different from a - S - S - group (disulfide bridge),  
 and acid addition salts thereof.

The abbreviations used in the formulae are in agreement with the nomenclature accepted in the peptide  
 chemistry, which has been published e.g. in J. Biol. Chem. 241, 527 (1966). According to this the  
 20 abbreviations occurring in the description are as follows:

	X	-X-
25	Phe phenylalanine	(phenylalanyl)
	Trp tryptophan	(tryptophyl)
	Tyr tyrosine	(tyrosyl)
	Sar sarcosine	(sarcosyl)
30	Ala alanine	(alanyl)
	His histidine	(histidyl)
	Val valine	(valyl)
35	Thr threonin	(threonyl)
	Pro proline	(prolyl)
	Leu leucine	(leucyl)
40	Cys cysteine	(cysteiny1)

45

50

55

	Phg	phenylglycine	(phenylglycyl)
	Pop	p-hydroxyphenylglycine	(p-hydroxyphenylglycyl)
5	Aa	o-aminobenzoic acid	(o-aminobenzoyl)
		anthranilic acid	(anthranilyl)
	Mab	m-aminobenzoic acid	(m-aminobenzoyl)
	Nal	$\beta$ -naphthyl-alanine	( $\beta$ -naphthyl-alanyl)
10	Tic	tetrahydroisoquinoline-carboxylic acid	(tetrahydroisoquinolyl-carbonyl)
	Asu	aminosuccinimide	(aminosuccinyl)
	Ac	acetyl	
15	Boc	tertiary butyloxycarbonyl	
	Bop	benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate	
	tBu	tertiary butyl	
20	Bzl	benzyl	
	DCC	dicyclohexylcarbodiimide	
	DCU	dicyclohexylurea	
	DIC	diisopropylcarbodiimide	
25	DIPEA	diisopropylethylamine	
	DMF	dimethylformamide	
	Et	ethyl	
30	Fmoc	(9-fluorenylmethyl)-oxycarbonyl	
	For	formyl	
	HPLC	high performance liquid chromatography	
	Me	methyl	
35	ONP	p-nitrophenyl	
	Opcp	pentachlorophenyl	
	Opfp	pentafluorophenyl	
	Ph	phenyl	
40	TEA	triethylamine	
	TEAA	triethylammonium acetate	
	TEAP	triethylammonium phosphate	
	TFA	trifluoroacetic acid	
45	Tfa	trifluoroacetyl	
	THF	tetrahydrofuran	
	TLC	thin layer chromatography	
	Tos	tosyl	
50	UV	ultraviolet	
	Z	benzyloxycarbonyl.	

Preferably the D-amino acid rest which may be represented by  $X_1$  is a D-phenylalanyl, D-naphthylalanyl, preferably D-2-naphthyl-alanyl, or D-tryptophyl rest, above all the first one, and/or the amino acid rest which may be represented by  $X_2$  is a tyrosyl rest and/or the halogen by which the rest of the D-amino acid derivative which may be represented by  $X_1$  and/or the rest of the amino acid derivative which may be represented by  $X_2$  may be ring-substituted is iodine, particularly preferably the rest of the D-amino acid




derivative which may be represented by  $X_1$  being a D-diiodotyrosyl rest and/or the rest of the amino acid derivative which may be represented by  $X_2$  being a diiodotyrosyl rest.

Furthermore it is preferred that the rest of the amine which may be represented by  $A_1$  is an indoliny1 or phenylamino rest, the alkoxy group which may be represented by  $A_1$  is such having from 1 to 4, particularly 1 or 2, carbon atom(s), the aryloxy group which may be represented by  $A_1$  is a phenyloxy group or the aralkyloxy group which may be represented by  $A_1$  is such having from 1 to 4, particularly 1 or 2, carbon atom(s) in the alkyl part and having as aryl part a phenyl part.

Moreover it is preferred that the alkyl side chain of the amino acid rest or its derivative which may be represented by  $X_4$  is such having from 1 to 4, particularly 3 or 4, carbon atom(s) or the aralkyl side chain or the amino acid rest or its derivative which may be represented by  $X_4$  is such having from 1 to 4, particularly 1 or 2, carbon atom(s) in the alkyl part and having as aryl part a phenyl part, particularly the amino acid rest which may be represented by  $X_4$  being a leucyl, valyl, isoleucyl, norvalyl, norleucyl, alanyl or seryl rest, above all one of the first two, and/or the aromatic amino acid rest which may be represented by  $X_5$  is a tryptophyl or phenylalanyl rest, above all the first one, or the halogen by which the rest of the derivative of the aromatic amino acid which may be represented by  $X_5$  may be substituted is iodine, particularly preferably the rest of the derivative of the aromatic amino acid which may be represented by  $X_5$  being a tyrosyl or diiodotyrosyl rest, and/or the  $C_{1-4}$  alkyl group which may be represented by  $A_2$  is such having 1 or 2 carbon atoms or the halogen by which the  $C_{1-4}$  alkyl group which may be represented by  $A_2$  may be substituted is fluorine the number of it being preferably 3, and/or  $n$  is 3 or 4, above all the latter one.

Particularly preferred compounds are



Suitably the acid addition salts of the octapeptide or heptapeptide derivatives according to the invention are such with pharmaceutically acceptable acids.

A subject matter of the invention is also a process for the preparation of the compounds according to the invention which is characterized in that using the method of the solid-phase peptide synthesis, the protected amino acids are stepwise condensed onto the solid resin carrier through carbodiimide or active ester coupling in the corresponding succession, then the final product is removed from the solid carrier by acidic or alkaline cleavage and the protective groups are removed from the amino acids simultaneously with or before or after cleavage from the resin and, if desired, the octapeptide or heptapeptide amide of the general formula I thus obtained is converted to an acid addition salt by reacting it with an acid and/or, if desired, the free base is liberated from the acid addition salt obtained by reacting it with an alkali and, if desired, the latter is converted to another acid addition salt by reacting it with an acid.

It is suitable to use a benzhydrylamine resin or a chloromethylated polystyrene resin as solid carrier in the process of the invention.

The final product containing protective groups can preferably be splitted off from the resin by using hydrogen fluoride and/or ammonolysis.

As already mentioned, the compounds of the invention have superior pharmacological activity, particularly in treating mammals, including man, suffering from tumour disease, and inhibiting tyrosine kinase activity.

Hence by the invention also there are provided for medicaments which are characterized in that they contain as active ingredient(s) 1 or more compound(s) according to the invention, suitably in admixture with 1 or more carrier(s) and/or additive(s) commonly used in the pharmaceutical technique.

The medicaments according to the invention can be in the form of pharmaceutical preparations. These preparations can be prepared by using methods known per se, and thus suitably by mixing 1 or more compound(s) according to the invention with 1 or more carrier(s) and/or additive(s) commonly used in the

pharmaceutical technique and transforming the mixture thus obtained to a pharmaceutical preparation.

The compounds according to the invention were found to be active in the dosage range of 0.01 to 500  $\mu\text{g/kg}$  of body-weight (hereinafter:  $\mu\text{g/kg}$ ) on mice or 0.5 to 2000 mg/kg on man, respectively.

Hence a further subject matter of the invention is the use of the compounds according to the invention for preparing medicaments, particularly for treating mammals, including man, suffering from tumour disease and/or for inhibiting their tyrosine kinase activity.

The pharmacological effects of the compounds according to the invention are supported by the tests described hereinafter.

#### 10 A) Assay of the growth hormone (GH)

The release or inhibition, respectively, of the release of GH were measured on rat hypophysis by using the superfusion method [Vigh et al.: Peptides 5, 241 (1984)]. As control, the GH amount was considered which was released by the GH releasing hormone (GHRH) given in the same dose as the sample.

15 The activity of the compound according to the invention was expressed as the percentage of decrease or increase, respectively, in the GH amount released by GHRH (Table I).

#### B) Assay of the inhibition of cell division

20 The incorporation of [ $^3\text{H}$ ]-thymidine to tumour cells of various origin and measurement of the cell count were carried out according to the method of Kéri et al. [Tumor Biology 9, 315 (1988)]. The biological activity of the compounds according to the invention was expressed as the percentage of inhibition of the labelled thymidine incorporation or increase in the count of untreated cells used as control (Table II).

#### 25 C) Measurement of the tyrosine kinase activity

The tyrosine kinase activity was also determined according to the method of Kéri et al. [Tumor Biology 9, 315 (1988)]. The activity of the compounds according to the invention was characterized on the basis of their inhibitory effect in comparison to the incorporation of  $^{32}\text{P}$  isotope to untreated cells used as control 30 (Table III).

#### D) Study of effectivity of the compounds on the tumour growth in the metastasis model

The anti-metastatic effect of the compounds was studied on Lewis lung tumour (LLT) cells in muscle-lung and spleen-liver metastasis model as well as on their immunoresistant cell variant (LLT-HH). These 35 experiments were carried out on inbred C<sub>57</sub>B1 mice of both sexes. The LLT was transplanted into the muscle for developing the muscle-lung metastasis model, whereas a suspension of the tumour cells was injected to the spleen to form the spleen-liver metastasis. The compounds to be tested were administered intraperitoneally (i.p.), intravenously (i.v.), orally (p.o.) or subcutaneously (s.c.) in 0.1 to 10 mg/kg doses 40 daily 1 to 3 times in the 5th to 13th days. The therapeutic effect was evaluated on the basis of the number of metastases in such a way that a sample was taken in the spleen-liver model between the 10th and 14th days, in the muscle-lung model between the 17th and 18th days and the macroscopic metastases in the liver and lungs, respectively, were counted under a stereomicroscope. The efficiency of the compounds was expressed in percentage of the count of metastases determined in the control animals (Table IV).

50

55

Table I

Change of GH release elicited by GHRH  
under effect of the analogues of the  
invention

5

10

15

20

25

30

35

40

45

50

55

Designation	Compound Example	Change of GH release
B-aspartyl-(indoliny1)- -Cys-Tyr-D-Trp-Lys-Val- -Cys-Thr-NH <sub>2</sub>	1	-100
D-Phe-Cys-Tyr-D-Trp- -Lys-Cys-Thr-NH <sub>2</sub>	2	0
D-Phe-Cys-Tyr-D-Trp- -Lys-Leu-Cys-Thr-NH <sub>2</sub>	3	-100
D-Phe-Cys-Tyr-B-Asp(in- doliny1)-Lys-Leu-Cys- -Thr-NH <sub>2</sub>	4	+290
D-Phe-Tyr-Tyr-Aa-Lys- -Val-Phe-Trp-NH <sub>2</sub>	5	+ 25
Ac-Sar-Ala-Tyr-Tyr-Aa- -Lys-Val-Phe-Trp-NH <sub>2</sub>	6	- 20
D-Tic-Cys-Tyr-D-Trp-Lys- -Val-Cys-Thr-NH <sub>2</sub>	10	- 85

Table I continued

5	Designation	Compound Example	Change of GH release
10	D-Phe-Cys-His-D-Trp-Lys- -Val-Cys-Thr-NH <sub>2</sub>	11	-100
15	D-Phe-Cys-Tyr-D-Trp-Lys- -B-Ala-Cys-Thr-NH <sub>2</sub>	13	- 10
20	D-Phe-Cys-Tyr-D-Trp-Lys- -Pro-Cys-Thr-NH <sub>2</sub>	14	0
25	Ac-Sar-Tyr-Tyr-Aa-Lys- -Val-Phe-Trp-NH <sub>2</sub>	15	+ 9

30

35

40

45

50

55

Table II

Assay of the inhibition of cell  
division on various tumour cells

5

10

15

20

25

30

35

40

45

50

55

Designation	Compound Example	Cell line <sup>+</sup>	Dose (µg)	Inhibition (%)
B-aspartyl-(indoliny1)- -Cys-Tyr-D-Trp-Lys-Val- -Cys-Thr-NH <sub>2</sub>	1	HT 29 HT 29 MCF 7	1 10 10	36 44 46
D-Phe-Cys-Tyr-D-Trp- -Lys-Cys-Thr-NH <sub>2</sub>	2	HT 29 DU 145 PC 3	20 10 10	75 68 45
D-Phe-Cys-Tyr-D-Trp- -Lys-Leu-Cys-Thr-NH <sub>2</sub>	3	HT 29 ET 29 MCF 7 DU 145	1 10 10 10	57 50 31 36
D-Phe-Cys-Tyr-β-Asp(in- doliny1)-Lys-Leu-Cys- -Thr-NH <sub>2</sub>	4	SW 620	10	68
D-Phe-Tyr-Tyr-Aa-Lys- -Val-Phe-Trp-NH <sub>2</sub>	5	MCF 7	40	48
Ac-Sar-Ala-Tyr-Tyr-Aa- -Lys-Val-Phe-Trp-NH <sub>2</sub>	6	HT 29 MCF 7	10 20	28 24
D-Tic-Cys-Tyr-D-Trp-Lys- -Val-Cys-Thr-NH <sub>2</sub>	10	MCF 7	10	21

Table II continu d

Designation	Compound Example	Cell line <sup>+</sup>	Dose ( $\mu$ S)	Inhibi- tion (%)
D-Phe-Cys-His-D-Trp- -Lys-Val-Cys-Thr-NH <sub>2</sub>	11	MCF 7 MCF 7	10 20	38 44
D-Phe-Cys-Tyr-D-Trp- -Lys(Tfa)-Val-Cys-Thr- -NH <sub>2</sub>	12	MCF 7 MCF 7	10 20	13 19
D-Phe-Cys-Tyr-D-Trp- -Lys- $\beta$ -Ala-Cys-Thr-NH <sub>2</sub>	13	MCF 7 HT 29	20 20	45 13

- <sup>+</sup> HT 29: tumour cell line of human colon origin  
 MCF 7: tumour cell line of human breast origin  
 DU 145: tumour cell line of human prostate  
 PC 3: tumour cell line of human prostate  
 SW 620: tumour cell line of human colon origin

Table III

Assay of th tyrosine kinase activity

5

10

15

20

25

30

35

40

45

50

55

Designation	Compound		Cell line <sup>+</sup>	Dose ( $\mu$ g)	Inhibi- tion (%)
	Example				
<u>B-aspartyl-(indoliny1)-</u> <u>-Cys-Tyr-D-Trp-Lys-Val-</u> <u>-Cys-Thr-NH<sub>2</sub></u>	1	HT 29	10	44	
<u>D-Phe-Cys-Tyr-D-Trp-</u> <u>-Lys-Cys-Thr-NH<sub>2</sub></u>	2	SW 620 SW 620	1 10	84 70	
<u>D-Phe-Cys-Tyr-D-Trp-</u> <u>-Lys-Leu-Cys-Thr-NH<sub>2</sub></u>	3	HT 29 HT 29	10 30	50 90	
<u>D-Phe-Cys-Tyr-B-Asp(in-</u> <u>doliny1)-Lys-Leu-Cys-</u> <u>-Thr-NH<sub>2</sub></u>	4	HT 29 HT 29	10 30	30 45	
<u>Ac-Sar-Ala-Tyr-Tyr-Aa-</u> <u>-Lys-Val-Phe-Trp-NH<sub>2</sub></u>	6	HT 29 HT 29	10 30	17 23	

<sup>+</sup> HT 29: tumor cell line of human colon origin  
SW 620: tumor cell line of human colon origin

Table IV

Effect on the tumour growth in the  
metastasis model

Compound Designation	Example	Count of metastases (%) related to controll (100%)
B-aspartyl-(indoliny)- -Cys-Tyr-D-Trp-Lys-Val- -Cys-Thr-NH <sub>2</sub>	1	43
D-Phe-Cys-Tyr-D-Trp- -Lys-Leu-Cys-Thr-NH <sub>2</sub>	3	35
D-Phe-Cys-Tyr-B-Asp(in- doliny)-Lys-Leu-Cys- -Thr-NH <sub>2</sub>	4	25
D-Phe-Cys-Tyr-D-Trp- -Lys(Tfa)-Val-Cys-Thr- -NH <sub>2</sub>	12	51

The main advantages of the invention are as follows:

- The compounds according to the invention can preferably be utilized to inhibit tumour growth or the activity of tyrosine kinase enzymes playing an important role in the tumorous transformation.
- The compounds according to the invention with the new amino acid combinations are useful also for regulating the release of growth hormone (GH), insulin, glucagon and prolactin and/or for inhibiting tumour growth.
- From the compounds according to the invention bearing an aminosuccinyl, o- and m-aminobenzoyl substitution, those containing no disulfide bridge do not exert any inhibitory effect on the GH release but inhibit the growth of tumour cells.
- The compounds according to the invention can be used to inhibit pathological processes, such as psoriasis, elicited by the pathological proliferation of skin cells.
- The o- and m-aminobenzoyl substituents are devoid of any centre of asymmetry, therefore no racemization can occur during their synthesis. Thus, the preparation of the final product becomes easier and cheaper.
- The preparation costs of o- and m-aminobenzoic acid are much lower than those of the D-amino acids playing a similar role in somatostatin analogues known up to now.

The invention is illustrated in detail by the following Examples.

In the Examples, the compounds are prepared by methods commonly used in the solid-phase peptide synthesis (Stewart et al.: Solid Phase Peptide Synthesis, 2nd Edition, Pierce Chemical Company, Rockford, Illinois, 1984). The individual amino acids are stepwise bound in the form of their Boc or Fmoc derivatives to a benzhydrylamine or chloromethylated polystyrene resin by the symmetric anhydride or active ester coupling or by the aid of DCC, DIC or BOP reagents.



The progress of the reaction is evaluated by the ninhydrin test [E. Kaiser et al.: Anal. Biochem. 34, 595 (1970)]. The acylation is repeated when a free amino group is detected. The time demand of coupling depends on the amino acids and varies between 1 and 16 hours.

Both the removal of protective groups and cleavage of the peptide from the resin are preferably carried out in a single step by using anhydrous liquid hydrogen fluoride [S. Sakakibara et al.: Bull. Chem. Soc. Japan 40, 2164 (1967)].

If desired, the crude product obtained by HF cleavage is cyclized to form the disulphide bridge.

The crude product is purified by using Sephadex chromatography and/or preparative HPLC method. The purity of the final product is examined by TLC, analytical HPLC and amino acid analysis. The TLC  $R_f$  values are determined on Kieselgel sheets (DC Alufolien, Merck) by using the solvent mixtures listed hereinafter the ratios being by volume:

15	1. Ethyl acetate/pyridine/acetic acid/water	30:20:6:11
	2. Ethyl acetate/pyridine/acetic acid/water	60:20:6:11
	3. Butanol/pyridine/acetic acid/water	60:20:6:11
	4. n-Butanol/acetic acid/water	4:1:2
	5. n-Butanol/acetic acid/water/ethyl acetate	1:1:1:1
	6. n-Butanol/acetic acid/water	4:1:1
20	7. Isopropanol/1 molar acetic acid	2:1
	8. Ethyl acetate/pyridine/acetic acid/water	5:5:1:3

In all Examples the percentages of solvents and eluents are by volume.

#### 25 Example 1

#### 30 Preparation of $\beta$ -aspartyl-(indolinyl)-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub> (Mw: 1076)

0.47 g (0.25 mmol) of benzhydrylamine hydrochloride resin (0.54 milliequivalent/g) is swollen in methylene chloride (CH<sub>2</sub>Cl<sub>2</sub>) for 90 minutes, then the following cycle is repeated after each acylation carried out with the amino acids. The percentages are by volume.

Step	Reagents, procedure	Time of stirring (min)
40	1 CH <sub>2</sub> Cl <sub>2</sub> ; 3 washings	1
	2 Mixture containing 33 % of TFA, 5 % of anisole and 62 % of CH <sub>2</sub> Cl <sub>2</sub> ; cleavage	1
	3 Mixture containing 33 % of TFA, 5 % of anisole and 62 % of CH <sub>2</sub> Cl <sub>2</sub> ; cleavage	1
45	4 CH <sub>2</sub> Cl <sub>2</sub> ; 3 washings	1
	5 Ethanol; 2 washings	1
	6 CH <sub>2</sub> Cl <sub>2</sub> ; 3 washings	1
	7 Mixture containing 10 % of TEA and 90 % of CH <sub>2</sub> Cl <sub>2</sub> ; 2 washings	2
	8 CH <sub>2</sub> Cl <sub>2</sub> ; 3 washings	2
	9 Ethanol; 2 washings	1
50	10 CH <sub>2</sub> Cl <sub>2</sub> ; 3 washings	1
	11 3 equivalents of Boc-amino acid dissolved in the mixture of CH <sub>2</sub> Cl <sub>2</sub> and DMF as well as 3 equivalents of DCC or DIC dissolved in CH <sub>2</sub> Cl <sub>2</sub> ; coupling	60 minutes to 16 hours
	12 CH <sub>2</sub> Cl <sub>2</sub> ; 3 washings	1
55	13 Ethanol; 2 washings	1

A ninhydrin test is made after each step. When the result is positive, the cycle is repeated from the 6th step.

0.25 mmol of  $\beta$ -Asp(indoliny)-Cys(4-Me-Bzl)-Tyr(2-Br-Z)-D-Trp-Lys(Z)-Val-Cys(4-Me-Bzl)-Thr(Bzl) peptide resin is suspended in 3 ml of anisole containing 10 mixed % of p-cresol, and 30 ml of HF gas are condensed onto the mixture. After stirring at 0 °C for 45 minutes HF is removed, the residue is suspended in 200 ml of ethyl acetate, stirred for 15 minutes, then poured into 250 ml of ethyl acetate. After filtration the precipitate is filtered, washed with ethyl acetate, then the peptide is dissolved from the precipitate by filtration, using 500ml of 95 % by volume gas-free acetic acid. Then 9.0 ml of 0.03 M iodine/methanol solution are dropped to the acetic acid solution until it becomes orange yellow. Thereafter, the solution is stirred for one hour. Zinc powder is cautiously added to the solution until the yellow colour disappears, then the zinc powder is filtered and the solution is evaporated. The solid residue is dissolved in 6 ml of 50 % by volume acetic acid and purified on a Sephadex® G-25 column. The elution is followed by using UV absorption measured at 280 nm and TLC. The fractions containing the aimed product are collected, evaporated, then further purified by gradient elution in a preparative HPLC system (column: Whatman Partisil 10 ODS-3 22x250 mm; eluent A: 0.25 N TEAP, pH 2.24; eluent B: 80 % of methanol with 20 % of A). The pure product is desalinated by gradient elution, using 0.02 M  $\text{NH}_4\text{OAc}$ , pH 5 (eluent A) as well as 70 % of methanol with 30 % of eluent A (eluent B) to obtain 120 mg (43 %) of final product. The physical characteristics of this compound are summarized in Table V, the biological effects already have been shown in Tables I, II, III and IV.

#### Example 2

#### Preparation of D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr- $\text{NH}_2$ (Mw: 947)

Starting from 0.47 g (0.25 mmol) of benzhydrylamine hydrochloride resin (0.54 milliequivalent/g) the peptide is synthesized, cyclized and purified as described in Example 1 to gave the final product in a yield of 80.5 mg (34 %). The physical characteristics are summarized in Table V, the biological effects already have been shown in Tables I and II.

#### Example 3

#### Preparation of D-Phe-Cys-Tyr-D-Trp-Lys-Leu-Cys-Thr- $\text{NH}_2$ (Mw: 1060)

Starting from 0.47 g (0.25 mmol) of benzhydrylamine hydrochloride resin (0.54 milliequivalent/g) the aimed peptide is synthesized, cyclized and purified as described in Example 1 to obtain 106 mg (40 %) of final product. The physical characteristics of this compound are summarized in Table V, the results of biological assays already have been shown in Tables II, III, IV and V.

#### Example 4

#### Preparation of D-Phe-Cys-Tyr- $\beta$ -Asp(indoliny)-Lys-Leu-Cys-Thr- $\text{NH}_2$ (Mw: 1076)

Starting from 0.47 g (0.25 mmol) of benzhydrylamine hydrochloride resin the aimed peptide is synthesized, cyclized and purified as described in Example 1 to give 94 mg (35.5 %) of final product. The physical characteristics of this compound are summarized in Table V, the results of biological assays are shown in Tables I, II, III and IV.

Example 5

Preparation of D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> (Mw: 1192)

- 5 Starting from 0.47 g (0.25 mmol) of benzhydrylamine hydrochloride resin the peptide is synthesized analogously as described in Example 1. The D-Phe-Tyr(2-Br-Z)-Tyr(2-Br-Z)-Aa-Lys(2-Cl-Z)-Val-Phe-Trp peptide resin obtained after the last step is suspended in 3 ml of anisole, 30 ml of gaseous HF are condensed onto the mixture which is then stirred at 0 °C for 45 minutes. After removing the gaseous HF the residue is suspended in 200 ml of abs. ether, stirred for 15 minutes and after filtration the precipitate is  
 10 washed with ether. Thereafter, the peptide is dissolved from the precipitate with 50 % by volume acetic acid, the solution is evaporated and the thus-obtained D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> heptapeptide amide is purified by preparative HPLC. Eluent A: 0.25 N TEAP, pH 2.25; eluent B: mixture of 70 % of acetonitrile and 30 % of eluent A. The separation is carried out on the column mentioned in Example 1 by using a mixture containing 30 % of eluent B and 70 % of eluent A. The pure product is desalinated as  
 15 described in Example 1 to obtain 89 mg (30 %) of final product. The physical characteristics of this compound are summarized in Table V, the biological effects already have been shown in Tables I and II.

Example 6

- 20 Preparation of Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> (Mw: 1217)

The aimed peptide is prepared by using the process of Example 5 but an Ac-Sar group is coupled to the peptide resin in the last cycle. The cleavage of the product from the resin and the purification are also carried out as described in Example 6 to obtain 85 mg (28 %) of final product. The physical characteristics  
 25 of the compound are summarized in Table V whereas its biological effects already have been shown in Tables I, II and III.

Example 7

- 30 Preparation of Pop-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> (Mw: 1194)

This peptide is synthesized as described in Example 1, except that in the 11th step Bop reagent is used in an amount equal to the amino acid as activating agent for coupling the amino acid in the presence of an excess of DIPEA. After cleavage by HE and purification by chromatography, the final product is obtained as  
 35 described in Example 5 with a yield of 105 mg (35 %). The physical characteristics of the compound are summarized in Table V.

Example 8

- 40 Preparation of Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub> (Mw: 1283)

After swelling 0.47 g (0.25 mmol) of benzhydrylamine hydrochloride resin in methylene chloride for 90 minutes, the following cycle is repeated after each acylation carried out with the Fmoc-derivatives of the suitably protected amino acids the percentages being by volume:  
 45

50

55

Step	Reagents, procedure	Time of stirring (minute)
1	DMF; 3 washings	2
2	Mixture containing 20 % of piperidine and 80 % of DMF; cleavage	2
3	Mixture containing 20 % of piperidine and 80 % of DMF; cleavage	10
4	DMF; 5 washings	2
5	Symmetric anhydride <sup>+</sup> prepared from 3 equivalents of Fmoc-amino acid or Opfp ester; coupling	60
6	DMF; 3 washings	2


<sup>+</sup> = Preparation of the symmetric anhydride: 3 equivalents of Fmoc-amino acid are dissolved in a mixture of CH<sub>2</sub>Cl<sub>2</sub> and DMF. After adding 1.5 equivalents of DCC the mixture is stirred at room temperature for 15 minutes. The DCU precipitate is filtered off, the solution is evaporated and the residue dissolved in DMF is used for coupling.

A ninhydrin test is made after each step. When a positive result is obtained, the cycle is repeated from the 5th step.

The Aa-Cys(Acm)-Tyr(tBu)-D-Trp-Lys(Boc)-Val-Cys(Acm)-Trp peptide resin obtained after the last step is treated with HF, purified and desalinized as described in Example 5 to obtain 147 mg (46 %) of final product. The physical characteristics of the compound are summarized in Table V.

#### Examples 9 to 14

The compounds of the following formulae, listed hereinafter, are prepared by using the process described in Examples 1 to 4.



Example	A	B	C	E
10	β-Asp(NH-Ph)	Tyr	Lys	Val
11	D-Tic	Tyr	Lys	Val
12	D-Phe	His	Lys	Val
13	D-Phe	Tyr	Lys(Tfa)	Val
14	D-Phe	Tyr	Lys	β-Ala
15	D-Phe	Tyr	Lys	Pro

The physical characteristics of the above compounds are summarized in Table V, the results of biological assays already have been shown in Tables I, II and V.

Examples 15 and 16

Compounds of the following formulae are prepared as described in Example 6:

5

A-B-Tyr-Aa-Lys-Val-Phe-Trp-NH <sub>2</sub>		
Example	A	B
15	Ac-Sar	Tyr
16	D-Phe	Ala


10

The physical characteristics of the above compounds are summarized in Table V, the results of biological assays already have been shown in Table I.

15 Examples 17 and 18

The following compounds are prepared by using the process described in Example 1:

20




In Example 17 "A" stands for a D-Phe group, in Example 18 for a D-2-naphthyl-alanyl group. The physical data of the compounds are summarized in Table V.

25

Example 19

The following compound is prepared by using the process described in Example 1:

30




The physical data of the compounds are summarized in Table V.

35

Example 20

40



The compound is prepared by using the process described in Example 1 with the exception that after coupling the Lys residue the neutralization is carried out with 5 % by weight diisopropyl-ethylamine in dichloromethane, instead of using TEA. The side-chain protecting group (fluorenylmethyl ester) of Asp is removed from the protected peptide resin by stirring it for 2x20 minutes in a 1:1 mixture of DMF and piperidine. The formation of Asp- $\beta$ -pentafluorophenyl ester is carried out by stirring the suspension of the peptide resin for 30 minutes in DMF with an excess of pentafluorophenol and diisopropyl carbodiimide. The peptide resin is filtered and washed with DMF. The aspartimide derivative is formed from the pentafluorophenyl ester derivative by stirring the peptide resin in DMF. The formation of the aspartimide ring is followed by measuring the released pentafluorophenol. The UV absorption of a small amount of the filtered reaction mixture is measured at 278 nm using DMF as control. The cleavage of the remaining protecting groups, the cleavage of the peptide from the resin, the formation of the disulfide bridge and the purification of the crude product are carried out using the process described in Example 1. The physical data of the compound are summarized in Table V.

Table V  
Physical characteristics of compounds of the invention

Compound		Example	M.p. (°C)	$[\alpha]_D^{20}$	R <sub>f</sub> <sup>1</sup>	R <sub>f</sub> <sup>2</sup>	R <sub>f</sub> <sup>3</sup>	R <sub>f</sub> <sup>4</sup>	R <sub>f</sub> <sup>5</sup>	R <sub>f</sub> <sup>6</sup>	R <sub>f</sub> <sup>7</sup>	R <sub>f</sub> <sup>8</sup>
β-aspartyl-(indolyl- -Cys-Tyr-D-Trp-Lys-Val- -Cys-Thr-NH <sub>2</sub> )		1		-53	0.62	0.19	0.61	0.55		0.56	0.81	
D-Phe-Cys-Tyr-D-Trp- -Lys-Cys-Thr-NH <sub>2</sub> )		2		-98.2	0.55			0.41		0.36	0.73	
D-Phe-Cys-Tyr-D-Trp- -Lys-Leu-Cys-Thr-NH <sub>2</sub> )		3		-43	0.65	0.22	0.66	0.59		0.61	0.81	
D-Phe-Cys-Tyr-β-Asp(in- dolyl)-Lys-Leu-Cys- -Thr-NH <sub>2</sub> )		4		-63	0.61	0.19	0.64	0.51		0.55	0.16	
D-Phe-Tyr-Tyr-Aa-Lys- -Val-Phe-Trp-NH <sub>2</sub> )		5	156-160	-19.6					0.83	0.69		0.88
Ac-Sar-Ala-Tyr-Tyr-Aa- -Lys-Val-Phe-Trp-NH <sub>2</sub> )		6	168-172	-34					0.70	0.60	0.70	0.60

Table V continued

Compound		M.p. (°C)	$[\alpha]_D^{20}$	$R_f^1$	$R_f^2$	$R_f^3$	$R_f^4$	$R_f^5$	$R_f^6$	$R_f^7$	$R_f^8$
Designation	Example										
Pop-Tyr-Tyr-As-Lys-Val-Phe-Trp-NH <sub>2</sub>	7							0.80	0.60		0.90
As-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH <sub>2</sub>	8	138-140	-47		0.27			0.84	0.40		0.81
$\beta$ -Aspartyl-(phenyl-amino)-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH <sub>2</sub>	9			0.63	0.26		0.72				
D-Tic-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH <sub>2</sub>	10		-57.4	0.63	0.22	0.64	0.51		0.55	0.80	
D-Phe-Cys-His-D-Trp-Lys-Val-Cys-Thr-NH <sub>2</sub>	11		-116.2	0.43	0.04	0.35	0.36		0.14	0.16	
D-Phe-Cys-Tyr-D-Trp-Lys(Trp)-Val-Cys-Thr-NH <sub>2</sub>	12		-68.6	0.85	0.63	0.76	0.75		0.78	0.93	
D-Phe-Cys-Tyr-D-Trp-Lys- $\beta$ -Ala-Cys-Thr-NH <sub>2</sub>	13		-30.8	0.60	0.01	0.54	0.48		0.43	0.80	

Table V continued

Compound		Example	M.p. (°C)	$[\alpha]_D^{20}$	R <sub>f</sub> <sup>1</sup>	R <sub>f</sub> <sup>2</sup>	R <sub>f</sub> <sup>3</sup>	R <sub>f</sub> <sup>4</sup>	R <sub>f</sub> <sup>5</sup>	R <sub>f</sub> <sup>6</sup>	R <sub>f</sub> <sup>7</sup>	R <sub>f</sub> <sup>8</sup>
Designation												
D-Phe-Cys-Tyr-D-Trp- Lys-Pro-Cys-Thr-NH <sub>2</sub>		14		-63.4	0.57	0.12	0.48	0.41		0.36	0.73	
Ac-Sar-Tyr-Tyr-Aa-Lys- Val-Phe-Trp-NH <sub>2</sub>		15			0.57	0.25	0.58			0.57		
D-Phe-Ala-Tyr-Aa-Lys- Val-Phe-Trp-NH <sub>2</sub>		16							0.80	0.60		0.90
D-Phe-Cys-Tyr-Aa-Lys- Val-Cys-Trp-NH <sub>2</sub>		17							0.61	0.38		
D-2-Naphthyl-Ala-Cys- Tyr-Aa-Lys-Val-Cys- Trp-NH <sub>2</sub>		18			0.09				0.74	0.49		
Aa-Cys-Tyr-D-Trp-Lys- Val-Cys-Trp-NH <sub>2</sub>		19			0.88					0.79		
D-Phe-Cys-Tyr-Aa-Lys- Val-Cys-Thr-NH <sub>2</sub>		20		-31.2		0.13	0.51	0.41				

\* c = 0,5 (0,1% by volume acetic acid)



## Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, PT, SE

1. Octapeptide or heptapeptide amide derivatives of the general formula

5

10

15

20

25

30


35

40

45

50

55




wherein

X<sub>1</sub> stands for an aromatic D-amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a D-phenylglycyl, p-hydroxyphenylglycyl, o-aminobenzoyl, m-aminoben-

zoyl, D-tetrahydroisoquinolylcarbonyl or sarcosylalanyl; or a

5




10

in which latter

- 15  $\text{A}_1$  means the rest of an aromatic, heterocyclic or aliphatic amine having a primary or secondary amino group  
 or means a primary or secondary aromatic, heterocyclic or aliphatic amino group or means an alkoxy, aryloxy or aralkyloxy group;  
 $\text{X}_2$  represents an aromatic amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a histidyl group;  
 20  $\text{X}_3$  means a D-tryptophyl, o-aminobenzoyl, m-aminobenzoyl, aspartyl, D-aspartyl,  $\beta$ -aspartyl,  $\beta$ -D-aspartyl or aminosuccinyl group; or a

25




30

in which latter

- 35  $\text{A}_1$  is as defined above;  
 $\text{X}_4$  stands for an amino acid rest bearing an alkyl or aralkyl side chain, or a derivative thereof substituted by a hydroxyl or methyl group in  $\beta$ -position; or a prolyl or  $\beta$ -alanyl group; or a valence bond;  
 $\text{X}_5$  means a heterocyclic amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a threonyl group;  
 40  $\text{R}_1$  represents hydrogen or a

45



in which latter

- 50  $\text{A}_2$  means hydrogen or a  $\text{C}_{1-4}$  alkyl group, optionally substituted by halogen;  
 $\text{R}_2$  means hydrogen, -S- or a -S-acetamidomethyl, phenyl or p-hydroxyphenyl group;  
 $\text{R}_3$  stands for hydrogen or hydroxyl;  
 $\text{R}_4$  means a phenyl or p-hydroxyphenyl group; or -S- or a -S-acetamidomethyl group when the meaning of  $\text{R}_2$  is the same group; or a methyl group when  $\text{R}_3$  is hydroxyl, in case of  $\text{R}_2$  and  $\text{R}_4$  each being -S- one bond each of them are united to a sole bond representing a bridge  
 55 between  $\text{R}_2$  and  $\text{R}_4$  and  
 $n$  is 1, 2, 3 or 4,  
 with th provisos that

- a) if
- X<sub>1</sub> stands for D-β-Nal or D-Phe, optionally substituted by halogen or a hydroxy group,
  - X<sub>2</sub> means Tyr or Phe
  - X<sub>3</sub> represents D-Trp,
  - X<sub>5</sub> stands for Trp or Thr and
  - R<sub>2</sub> and R<sub>4</sub> each are -S-, one bond each of them being united to a sole bond representing a bridge between R<sub>2</sub> and R<sub>4</sub>
  - X<sub>4</sub> is different from Val, Thr or Ser,
- b) if
- X<sub>4</sub> stands for Val or Thr or Ser and
  - X<sub>1</sub> means an aromatic D-amino acid residue or a derivative thereof ring-substituted by halogen or hydroxy
  - X<sub>2</sub> represents an aromatic amino acid rest or a derivative ring-substituted by halogen or hydroxy
- at least one of the following conditions is fulfilled
- α)
- X<sub>3</sub> is different from D-Trp
- or
- β)
- R<sub>2</sub> and R<sub>4</sub> together have a meaning different from a - S - S - group (disulfide bridge), and acid addition salts thereof.
2. Octapeptide or heptapeptide amide derivatives according to claim 1, characterized in that the D-amino acid rest which may be represented by X<sub>1</sub> is a D-phenylalanyl, D-naphthylalanyl or D-tryptophyl rest and/or the amino acid rest which may be represented by X<sub>2</sub> is a tyrosyl rest and/or the halogen by which the rest of the D-amino acid derivative which may be represented by X<sub>1</sub> and/or the rest of the amino acid derivative which may be represented by X<sub>2</sub> may be ring-substituted is iodine, particularly preferably the rest of the D-amino acid derivative which may be represented by X<sub>1</sub> being a D-diiodotyrosyl rest and/or the rest of the amino acid derivative which may be represented by X<sub>2</sub> being a diiodotyrosyl rest.
3. Octapeptide or heptapeptide amide derivatives according to claim 1, characterized in that the rest of the amine which may be represented by A<sub>1</sub> is an indolyl or phenylamino rest, the alkoxy group which may be represented by A<sub>1</sub> is such having from 1 to 4 carbon atom(s), the aryloxy group which may be represented by A<sub>1</sub> is a phenyloxy group or the aralkyloxy group which may be represented by A<sub>1</sub> is such having from 1 to 4 carbon atom(s) in the alkyl part and having as aryl part a phenyl part.
4. Octapeptide or heptapeptide amide derivatives according to claims 1 to 3, characterized in that the alkyl side chain of the amino acid rest or its derivative which may be represented by X<sub>4</sub> is such having from 1 to 4 carbon atom(s) or the aralkyl side chain of the amino acid rest or its derivative which may be represented by X<sub>4</sub> is such having from 1 to 4 carbon atom(s) in the alkyl part and having as aryl part a phenyl part, particularly the amino acid rest which may be represented by X<sub>4</sub> being a leucyl, valyl, isoleucyl, norvalyl, norleucyl, alanyl or seryl rest and/or the aromatic amino acid rest which may be represented by X<sub>5</sub> is a tryptophyl or phenylalanyl rest, or the halogen by which the rest of the derivative of the aromatic amino acid which may be represented by X<sub>5</sub> may be substituted is iodine, particularly preferably the rest of the derivative of the aromatic amino acid which may be represented by X<sub>5</sub> being a tyrosyl or diiodotyrosyl rest, and/or the C<sub>1-4</sub> alkyl group which may be represented by A<sub>2</sub> is such having 1 or 2 carbon atoms or the halogen by which the C<sub>1-4</sub> alkyl group which may be represented by A<sub>2</sub> may be substituted is fluorine and/or n is 3 or 4.

5. A compound as claimed in claims 1 to 4, selected from the group consisting of

$\beta$ -aspartyl-(indoliny1)-Cys-

-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub>,

D-Phe-Cys-Tyr-D-Trp-Lys-

-Cys-Thr-NH<sub>2</sub>, D-Phe-Cys-Tyr-D-Trp-Lys-Leu-Cys-Thr-

-NH<sub>2</sub>, D-Phe-Cys-Tyr- $\beta$ -Asp(indoliny1)-Lys-Leu-Cys-

-Thr-NH<sub>2</sub>,

D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>, Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>, Pop-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> and Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub>.

6. A process for the preparation of the compounds according to claims 1 to 5, characterized in that using the method of the solid-phase peptide synthesis, the protected amino acids are stepwise condensed onto the solid resin carrier through carbodiimide or active ester coupling in the corresponding succession, then the final product is removed from the solid carrier by acidic or alkaline cleavage and the protective groups are removed from the amino acids simultaneously with or before or after cleavage from the resin and, if desired, the octapeptide or heptapeptide amide of the general formula I thus obtained is converted to an acid addition salt by reacting it with an acid and/or, if desired, the free base is liberated from the acid addition salt obtained by reacting it with an alkali and, if desired, the latter is converted to another acid addition salt by reacting it with an acid.
7. A process as claimed in claim 6, characterized in that a benzhydrylamine resin or a chloromethylated polystyrene resin is used as solid resin carrier.
8. A process as claimed in claim 6 or 7, characterized in that the final product containing protective groups is splitted off by hydrogen fluoride and/or ammonolysis.
9. Medicaments characterized in that they contain as active ingredient(s) 1 or more compound(s) as claimed in claims 1 to 8, suitably in admixture with 1 or more carrier(s) and/or additive(s) commonly used in the pharmaceutical technique.
10. The use of the compounds according to claims 1 to 5 for preparing medicaments, particularly for treating mammals, including man, suffering from tumour disease and/or for inhibiting their tyrosine kinase activity.

Claims for the following Contracting States : ES, GR

1. A process for the preparation of octapeptide or heptapeptide amide derivatives of the general formula

5

I,

10

15

20

25


30

35

40

45

50




55

wherein

$\text{X}_1$  stands for an aromatic D-amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a D-phenylglycyl, p-hydroxyphenylglycyl, o-aminobenzoyl, m-aminobenzoyl, D-tetrahydroisoquinolylcarbonyl or sarcosylalanyl; or a

5




10

in which latter

- A<sub>1</sub> means the rest of an aromatic, heterocyclic or aliphatic amine having a primary or secondary amino group  
 or means a primary or secondary aromatic, heterocyclic or aliphatic amino group or means an alkoxy, aryloxy or aralkyloxy group;
- X<sub>2</sub> represents an aromatic amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a histidyl group;
- X<sub>3</sub> means a D-tryptophyl, o-aminobenzoyl, m-aminobenzoyl, aspartyl, D-aspartyl, β-aspartyl, β-D-aspartyl or aminosuccinyl group; or a

20

25




30

in which latter

- A<sub>1</sub> is as defined above;
- X<sub>4</sub> stands for an amino acid rest bearing an alkyl or aralkyl side chain, or a derivative thereof substituted by a hydroxyl or methyl group in β-position; or a prolyl or β-alanyl group; or a valence bond;
- X<sub>5</sub> means an aromatic amino acid rest or a derivative thereof ring-substituted by halogen and/or hydroxyl; or a threonyl group;
- R<sub>1</sub> represents hydrogen or a

40



45

in which latter

- A<sub>2</sub> means hydrogen or a C<sub>1-4</sub> alkyl group, optionally substituted by halogen;
- R<sub>2</sub> means hydrogen, -S- or a -S-acetamidomethyl, phenyl or p-hydroxyphenyl group;
- R<sub>3</sub> stands for hydrogen or hydroxyl;
- R<sub>4</sub> means a phenyl or p-hydroxyphenyl group; or -S- or a -S-acetamidomethyl group when the meaning of R<sub>2</sub> is the same group; or a methyl group when R<sub>3</sub> is hydroxyl, in case of R<sub>2</sub> and R<sub>4</sub> each being -S- one bond each of them are united to a sole bond representing a bridge between R<sub>2</sub> and R<sub>4</sub>
- and

55

- n is 1, 2, 3 or 4,  
 with the provisos that

a) if

- X<sub>1</sub> stands for D-β-Nal or D-Phe, optionally substituted halogen or a hydroxy group,

X<sub>2</sub> means Tyr or Phe  
 X<sub>3</sub> represents D-Trp,  
 X<sub>5</sub> stands for Trp or Thr and  
 R<sub>2</sub> and R<sub>4</sub> each are -S-, one bond each of them being united to a sole bond representing a  
 5 bridge between R<sub>2</sub> and R<sub>4</sub>  
 X<sub>4</sub> is different from Val, Thr or Ser,

b) if  
 X<sub>4</sub> stands for Val or Thr or Ser and  
 X<sub>1</sub> means an aromatic D-amino acid residue or a derivative thereof ring-substituted by  
 10 halogen or hydroxy  
 X<sub>2</sub> represents an aromatic amino acid rest or a derivative ring-substituted by halogen or  
 hydroxy

at least one of the following conditions is fulfilled

α)  
 15 X<sub>3</sub> is different from D-Trp  
 or  
 β)

R<sub>2</sub> and R<sub>4</sub> together have a meaning different from a - S - S - group (disulfide bridge),  
 and acid addition salts thereof. characterized in that using the method of the solid-phase peptide  
 20 synthesis, the protected amino acids are stepwise condensed onto the solid resin carrier through  
 carbodiimide or active ester coupling in the corresponding succession, then the final product is  
 removed from the solid carrier by acidic or alkaline cleavage and the protective groups are removed  
 from the amino acids simultaneously with or before or after cleavage from the resin and, if desired, the  
 25 octapeptide or heptapeptide amide of the general formula I thus obtained is converted to an acid  
 addition salt by reacting it with an acid and/or, if desired, the free base is liberated from the acid  
 addition salt obtained by reacting it with an alkali and, if desired, the latter is converted to another acid  
 addition salt by reacting it with an acid.

2. A process as claimed in claim 1, characterized in that a benzhydrylamine resin or a chloromethylated  
 30 polystyrene resin is used as solid resin carrier.

3. A process as claimed in claim 1 or 2, characterized in that the final product containing protective  
 groups is splitted off by hydrogen fluoride and/or ammonolysis.


35 4. A process as claimed in claims 1 to 3, characterized in that octapeptide or heptapeptide amide  
 derivatives are prepared in which the D-amino acid rest which may be represented by X<sub>1</sub> is a D-  
 phenylalanyl, D-naphthylalanyl or D-tryptophyl rest and/or the amino acid rest which may be repre-  
 sented by X<sub>2</sub> is a tyrosyl rest and/or the halogen by which the rest of the D-amino acid derivative which  
 may be represented by X<sub>1</sub> and/or the rest of the amino acid derivative which may be represented by  
 40 X<sub>2</sub> may be ring-substituted is iodine, particularly preferably the rest of the D-amino acid derivative  
 which may be represented by X<sub>1</sub> being a D-diiodotyrosyl rest and/or the rest of the amino acid derivative  
 which may be represented by X<sub>2</sub> being a diiodotyrosyl rest.

45 5. A process as claimed in claims 1 to 3, characterized in that octapeptide or heptapeptide amide  
 derivatives are prepared in which the rest of the amine which may be represented by A<sub>1</sub> is an indoliny  
 or phenylamino rest, the alkoxy group which may be represented by A<sub>1</sub> is such having from 1 to 4  
 carbon atom(s), the aryloxy group which may be represented by A<sub>1</sub> is a phenyloxy group or the  
 aralkyloxy group which may be represented by A<sub>1</sub> is such having from 1 to 4 carbon atom(s) in the  
 50 alkyl part and having as aryl part a phenyl part.

6. A process as claimed in claims 1 to 3, characterized in that octapeptide or heptapeptide amide  
 derivatives are prepared in which the alkyl side chain of the amino acid rest or its derivative which may  
 be represented by X<sub>4</sub> is such having from 1 to 4 carbon atom(s) or the aralkyl side chain of the amino  
 acid rest or its derivative which may be represented by X<sub>4</sub> is such having from 1 to 4 carbon atom(s) in  
 55 the alkyl part and having as aryl part a phenyl part, particularly the amino acid rest which may be  
 represented by X<sub>4</sub> being a leucyl, valyl, isoleucyl, norvalyl, norleucyl, alanyl or seryl rest and/or the  
 aromatic amino acid rest which may be represented by X<sub>5</sub> is a tryptophyl or phenylalanyl rest, or the  
 halogen by which the rest of the derivative of the aromatic amino acid which may be represented by X<sub>5</sub>

may be substituted is iodine, particularly preferably the rest of the derivative or the aromatic amino acid which may be represented by  $X_5$  being a tyrosyl or diiodotyrosyl rest, and/or the  $C_{1-4}$  alkyl group which may be represented by  $A_2$  is such having 1 or 2 carbon atoms or the halogen by which the  $C_{1-4}$  alkyl group which may be represented by  $A_2$  may be substituted is fluorine and/or  $n$  is 3 or 4.

7. A process as claimed in claims 1 to 3, characterized in that octapeptide or heptapeptide amide derivatives are prepared which are selected from the group consisting of



D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>, Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>, Pop-Tyr-Tyr-Aa-  
 Lys-Val-Phe-Trp-NH<sub>2</sub>, Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub>.

8. A process for preparing medicaments characterized by mixing as active ingredient(s) 1 or more compound(s) prepared by the process according to claims 1 to 7, with 1 or more carrier(s) and/or additive(s) commonly used in the pharmaceutical technique in a manner known per se.



## Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, PT, SE

1. Octapeptid- oder Heptapeptidamidderivate der allgemeinen Formel

5

10

15

20

25

30


35

40

45

50


55



worin  
X<sub>1</sub>


für einen aromatischen D-Aminosäurerest oder ein Derivat davon, ringssubstituiert durch Halogen und/oder Hydroxyl; oder für ein D-Phenylglycyl, p-Hydroxyphenyl-glycyl, o-Amin-

obenzoyl, m-Aminobenzoyl, D-Tetrahydroisochinolylcarbonyl oder Sarcosylalanyl; oder für eine




steht, wobei in letzterer

- A<sub>1</sub> den Rest eines aromatischen, heterocyclischen oder aliphatischen Amins mit einer primären oder sekundären Aminogruppe, oder eine primäre oder sekundäre, aromatische, heterocyclische oder aliphatische Aminogruppe, oder eine Alkoxy-, Aryloxy- oder Aralkyloxygruppe bedeutet;
- X<sub>2</sub> für einen aromatischen Aminosäurerest oder ein Derivat davon, ringsubstituiert durch Halogen und/oder Hydroxyl; oder für eine Histidylgruppe steht;
- X<sub>3</sub> für eine D-Tryptophyl-, o-Aminobenzoyl-, m-Aminobenzoyl-, Aspartyl-, D-Aspartyl-, D-Aspartyl-, β-Aspartyl-, β-D-Aspartyl- oder Aminosuccinylgruppe; oder für eine



steht, wobei in letzterer

- A<sub>1</sub> die obenstehende Bedeutung besitzt;
- X<sub>4</sub> für einen eine Alkyl- oder Aralkylseitenkette tragenden Aminosäurerest oder ein Derivat davon, substituiert durch eine Hydroxyl- oder Methylgruppe in β-Position; oder für eine Prolyl- oder β-Alanylgruppe; oder für eine Valenzbindung steht;
- X<sub>5</sub> einen heterocyclischen Aminosäurerest oder ein Derivat davon, ringsubstituiert durch Halogen und/oder Hydroxyl; oder eine Threonylgruppe bedeutet;
- R<sub>1</sub> für Wasserstoff oder eine



steht,

wobei in letzterer

- A<sub>2</sub> für Wasserstoff oder eine C<sub>1-4</sub>-Alkylgruppe, wahlweise durch Halogen substituiert, steht;
- R<sub>2</sub> Wasserstoff, -S- oder eine -S-Acetamidomethyl-, Phenyl- oder p-Hydroxyphenylgruppe bedeutet;
- R<sub>3</sub> für Wasserstoff oder Hydroxyl steht;
- R<sub>4</sub> eine Phenyl- oder p-Hydroxyphenylgruppe; oder eine -S-Gruppe oder eine -S-Acetamidomethylgruppe, wenn R<sub>2</sub> die gleiche Gruppe bedeutet; oder eine Methylgruppe, wenn R<sub>3</sub> Hydroxyl ist, bedeutet, wobei im Falle, daß R<sub>2</sub> und R<sub>4</sub> jeweils eine -S-Bindung bedeuten, jeweils eine Bindung von diesen zu einer Bindung verbunden ist, was eine Brücke zwischen R<sub>2</sub> und R<sub>4</sub> ergibt


und worin

n 1, 2, 3 oder 4 ist,

mit der Maßgabe, daß wenn

- a)
- X<sub>1</sub> für D-β-Nal oder D-Phe, wahlweise durch Halogen oder eine Hydroxygruppe substituiert, steht,
- X<sub>2</sub> Tyr oder Phe bedeutet,
- 5 X<sub>3</sub> D-Trp bedeutet,
- X<sub>5</sub> Trp oder Thr bedeutet und
- R<sub>2</sub> und R<sub>4</sub> jeweils -S- bedeuten, wobei jeweils eine Bindung von diesen zu einer Bindung verbunden sind, was eine Brücke zwischen R<sub>2</sub> und R<sub>4</sub> ergibt, nicht Val, Thr oder Ser ist;
- 10 X<sub>4</sub>
- b)
- X<sub>4</sub> Val oder Thr oder Ser bedeutet und
- X<sub>1</sub> einen aromatischen D-Aminosäurerest oder ein Derivat davon, ringsubstituiert durch Halogen oder Hydroxy, bedeutet,
- X<sub>2</sub> für einen aromatischen Aminosäurerest oder ein Derivat, ringsubstituiert durch Halogen oder Hydroxy, steht,
- 15 mindestens eine der folgenden Bedingungen erfüllt ist:
- α)
- X<sub>3</sub> nicht D-Trp ist
- oder
- 20 β)
- R<sub>2</sub> und R<sub>4</sub> zusammen nicht die Bedeutung einer -S-S-Gruppe (Disulfidbrücke) besitzen; und Säureadditionssalze davon.
2. Octapeptid- oder Heptapeptidamidderivate gemäß Anspruch 1, **dadurch gekennzeichnet**, daß der D-Aminosäurerest, welcher durch X<sub>1</sub> repräsentiert werden kann, ein D-Phenylalanyl-, D-Naphthylalanyl- oder D-Tryptophylrest ist, und/oder der Aminosäurerest, der durch X<sub>2</sub> repräsentiert werden kann, ein Tyrosylrest ist, und/oder das Halogen, durch das der Rest des D-Aminosäurederivates, welches durch X<sub>1</sub> repräsentiert werden kann, und/oder der Rest des Aminosäurederivates, welcher durch X<sub>2</sub> repräsentiert werden kann, ring-substituiert werden kann, Iod ist, wobei besonders bevorzugt der Rest des D-Aminosäurederivates, welcher durch X<sub>1</sub> repräsentiert werden kann, ein D-Diiodtyrosylrest ist, und/oder der Rest des Aminosäurederivates, welcher durch X<sub>2</sub> repräsentiert werden kann, ein Diiodtyrosylrest ist.
3. Octapeptid- oder Heptapeptidamidderivate gemäß Anspruch 1, **dadurch gekennzeichnet**, daß der Rest des Amins, welches durch A<sub>1</sub> repräsentiert werden kann, ein Indoliny- oder Phenylaminorest ist, die Alkoxygruppe, welche durch A<sub>1</sub> repräsentiert werden kann, eine solche mit 1 bis 4 Kohlenstoffatomen ist, die Aryloxygruppe, welche durch A<sub>1</sub> repräsentiert werden kann, eine Phenylloxygruppe ist, oder die Aryloxygruppe, welche durch A<sub>1</sub> repräsentiert werden kann, eine solche mit 1 bis 4 Kohlenstoffatomen in dem Alkylteil und einem Phenylteil als Arylteil ist.
4. Octapeptid- oder Heptapeptidamidderivate gemäß Anspruch 1 bis 3, **dadurch gekennzeichnet**, daß die Alkylseitenkette des Aminosäurerestes oder seines Derivates, welcher durch X<sub>4</sub> repräsentiert werden kann, ein solcher mit 1 bis 4 Kohlenstoffatomen ist, oder die Arylseitenkette des Aminosäurerestes oder seines Derivates, welcher durch X<sub>4</sub> repräsentiert werden kann, eine solche mit 1 bis 4 Kohlenstoffatomen im Alkylteil und einem Phenylteil als Arylteil ist, wobei insbesondere der Aminosäurerest, welcher durch X<sub>4</sub> repräsentiert werden kann, ein Leucyl-, Valyl-, Isoleucyl-, Norvalyl-, Norleucyl-, Alanyl- oder Serylrest ist, und/oder der aromatische Aminosäurerest, welcher durch X<sub>5</sub> repräsentiert werden kann, ein Tryptophyl- oder Phenylalanylrest ist, oder das Halogen, durch das der Rest des Derivats der aromatischen Aminosäure, welche durch X<sub>5</sub> repräsentiert werden kann, substituiert werden kann, Iod ist, wobei besonders bevorzugt der Rest des Derivats der aromatischen Aminosäure, welche durch X<sub>5</sub> repräsentiert werden kann, ein Tyrosyl- oder Diiodtyrosylrest ist, und/oder die C<sub>1-4</sub>-Alkylgruppe, welche durch A<sub>2</sub> repräsentiert werden kann, eine solche mit 1 oder 2 Kohlenstoffatomen ist, oder das Halogen, durch das die C<sub>1-4</sub>-Alkylgruppe, welche durch A<sub>2</sub> repräsentiert werden kann, substituiert werden kann, Fluor ist und/oder n 3 oder 4 ist.

5. Verbindung gemäß Anspruch 1 bis 4, gewählt aus der aus



10

D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>,  
 Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>,  
 Pop-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> und Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub> bestehen-  
 den Gruppe.

15

6. Verfahren zur Herstellung der Verbindungen gemäß Anspruch 1 bis 5, **dadurch gekennzeichnet**, daß das Verfahren der Festphasen-Peptidsynthese angewandt wird, wobei die geschützten Aminosäuren schrittweise auf dem festen Harzträger durch Carbodiimidkopplung oder durch Kopplung mittels aktivem Ester in der entsprechenden Reihenfolge kondensiert werden, dann das Endprodukt von dem festen Träger durch saure oder alkalische Spaltung entfernt wird und die Schutzgruppen von den Aminosäuren gleichzeitig mit oder vor oder nach der Abspaltung vom Harz entfernt werden, und, falls gewünscht, das derart erhaltene Octapeptid- oder Heptapeptidamid der allgemeinen Formel I zu einem Säureadditionssalz durch Umsetzung mit einer Säure umgewandelt wird, und/oder, falls gewünscht, die freie Base von dem erhaltenen Säureadditionssalz durch Umsetzung mit einer Alkalie freigesetzt wird, und, falls gewünscht, letztere zu einem anderen Säureadditionssalz durch Umsetzung dieser mit einer Säure umgewandelt wird.
7. Verfahren gemäß Anspruch 6, **dadurch gekennzeichnet**, daß ein Benzhydrylaminharz oder ein chlormethyliertes Polystyrolharz als fester Harzträger verwendet wird.
8. Verfahren gemäß Anspruch 6 oder 7, **dadurch gekennzeichnet**, daß das Schutzgruppen enthaltende Endprodukt durch Fluorwasserstoff und/oder Ammonolyse aufgepalten wird.
9. Medikamente, **dadurch gekennzeichnet**, daß sie als aktive Bestandteile 1 oder mehrere Verbindungen gemäß den Ansprüchen 1 bis 8 enthalten, geeigneterweise in Vermischung mit 1 oder mehreren Trägern und/oder Additiven, wie sie üblicherweise in der pharmazeutischen Technik verwendet werden.
10. Verwendung der Verbindungen gemäß den Ansprüchen 1 bis 5 zur Herstellung von Medikamenten, insbesondere zur Behandlung von Säugetieren, einschließlich Menschen, die an Tumorerkrankungen leiden, und/oder zur Inhibierung ihrer Tyrosinkinaseaktivität.

45

50

55

Patentansprüche für folgende Vertragsstaaten : ES, GR

1. Verfahren zur Herstellung von Octapeptid- oder Heptapeptidamidderivaten der allgemeinen Formel

5

10

15

20

25

30

35

40

45

50

55


worin

 $X_1$ 

für einen aromatischen D-Aminosäurerest oder ein Derivat davon, ringssubstituiert durch Halogen und/oder Hydroxyl; oder für ein D-Phenylglycyl, p-Hydroxyphenyl-glycyl, o-Aminobenzoyl, m-Aminobenzoyl, D-Tetrahydroisochinoly carbonyl oder Sarcosylalanyl; oder für

eine

5




10

steht, wobei in letzterer

- A<sub>1</sub> den Rest eines aromatischen, heterocyclischen oder aliphatischen Amins mit einer primären oder sekundären Aminogruppe, oder eine primäre oder sekundäre, aromatische, heterocyclische oder aliphatische Aminogruppe, oder eine Alkoxy-, Aryloxy- oder Aralkyloxygruppe bedeutet;  
 X<sub>2</sub> für einen aromatischen Aminosäurerest oder ein Derivat davon, ringsubstituiert durch Halogen und/oder Hydroxyl; oder für eine Histidylgruppe steht;  
 X<sub>3</sub> für eine D-Tryptophyl-, o-Aminobenzoyl-, m-Aminobenzoyl-, Aspartyl-, D-Aspartyl-, D-Aspartyl-, β-Aspartyl-, β-D-Aspartyl- oder Aminosuccinylgruppe; oder für eine

20




25

steht, wobei in letzterer

- A<sub>1</sub> die obenstehende Bedeutung besitzt;  
 X<sub>4</sub> für einen eine Alkyl- oder Aralkylseitenkette tragenden Aminosäurerest oder ein Derivat davon, substituiert durch eine Hydroxyl- oder Methylgruppe in β-Position; oder für eine Prolyl- oder β-Alanylgruppe; oder für eine Valenzbindung steht;  
 X<sub>5</sub> einen heterocyclischen Aminosäurerest oder ein Derivat davon, ringsubstituiert durch Halogen und/oder Hydroxyl; oder eine Threonylgruppe bedeutet;  
 R<sub>1</sub> für Wasserstoff oder eine

35



40

steht,

wobei in letzterer

- A<sub>2</sub> für Wasserstoff oder eine C<sub>1</sub>-4-Alkylgruppe, wahlweise durch Halogen substituiert, steht;  
 R<sub>2</sub> Wasserstoff, -S- oder eine -S-Acetamidomethyl-, Phenyl- oder p-Hydroxyphenylgruppe bedeutet;  
 R<sub>3</sub> für Wasserstoff oder Hydroxyl steht;  
 R<sub>4</sub> eine Phenyl- oder p-Hydroxyphenylgruppe; oder eine -S-Gruppe oder eine -S-Acetamidomethylgruppe, wenn R<sub>2</sub> die gleiche Gruppe bedeutet; oder eine Methylgruppe, wenn R<sub>3</sub> Hydroxyl ist, bedeutet, wobei im Falle, daß R<sub>2</sub> und R<sub>4</sub> jeweils eine -S-Bindung bedeuten, jeweils eine Bindung von diesen zu einer Bindung verbunden ist, was eine Brücke zwischen R<sub>2</sub> und R<sub>4</sub> ergibt und worin

50

- n 1, 2, 3 oder 4 ist,  
 mit der Maßgabe, daß wenn

55


a)

- X<sub>1</sub> für D-β-Nal oder D-Phe, wahlweise durch Halogen oder eine Hydroxygruppe

- substituiert, steht,  
 X<sub>2</sub> Tyr oder Phe bedeutet,  
 X<sub>3</sub> D-Trp bedeutet,  
 X<sub>5</sub> Trp oder Thr bedeutet und  
 5 R<sub>2</sub> und R<sub>4</sub> jeweils -S- bedeuten, wobei jeweils eine Bindung von diesen zu einer Bindung verbunden sind, was eine Brücke zwischen R<sub>2</sub> und R<sub>4</sub> ergibt,  
 X<sub>4</sub> nicht Val, Thr oder Ser ist;  
 b)  
 X<sub>4</sub> Val oder Thr oder Ser bedeutet und  
 10 X<sub>1</sub> einen aromatischen D-Aminosäurerest oder ein Derivat davon, ringsubstituiert durch Halogen oder Hydroxy, bedeutet,  
 X<sub>2</sub> für einen aromatischen Aminosäurerest oder ein Derivat, ringsubstituiert durch Halogen oder Hydroxy, steht,  
 mindestens eine der folgenden Bedingungen erfüllt ist:  
 15 α)  
 X<sub>3</sub> nicht D-Trp ist  
 oder  
 β)  
 R<sub>2</sub> und R<sub>4</sub> zusammen nicht die Bedeutung einer -S-S-Gruppe (Disulfidbrücke) besitzen;  
 20 und Säureadditionssalze davon;  
**dadurch gekennzeichnet**, daß das Verfahren der Festphasen-Peptidsynthese angewandt wird, wobei die geschützten Aminosäuren schrittweise auf dem festen Harzträger durch Carbodiimidkopplung oder durch Kopplung mittels aktivem Ester in der entsprechenden Reihenfolge kondensiert werden, dann das Endprodukt von dem festen Träger durch saure oder alkalische Spaltung entfernt wird und die  
 25 Schutzgruppen von den Aminosäuren gleichzeitig mit oder vor oder nach der Abspaltung vom Harz entfernt werden, und, falls gewünscht, das derart erhaltene Octapeptid- oder Heptapeptidamid der allgemeinen Formel I zu einem Säureadditionssalz durch Umsetzung mit einer Säure umgewandelt wird, und/oder, falls gewünscht, die freie Base von dem erhaltenen Säureadditionssalz durch Umsetzung mit einer Alkalie freigesetzt wird, und, falls gewünscht, letztere zu einem anderen Säureadditionssalz durch Umsetzung dieser mit einer Säure umgewandelt wird.  
 30
2. Verfahren gemäß Anspruch 1, **dadurch gekennzeichnet**, daß ein Benzhydrylaminharz oder ein chlormethyliertes Polystyrolharz als fester Harzträger verwendet wird.
- 35 3. Verfahren gemäß Anspruch 1 oder 2, **dadurch gekennzeichnet**, daß das Schutzgruppen enthaltende Endprodukt durch Fluorwasserstoff und/oder Ammonolyse aufgepalten wird.
- 40 4. Verfahren gemäß den Ansprüchen 1 bis 3, **dadurch gekennzeichnet**, daß Octapeptid- oder Heptapeptidamidderivate hergestellt werden, in denen der D-Aminosäurerest, welcher durch X<sub>1</sub> repräsentiert werden kann, ein D-Phenylalanyl-, D-Naphthylalanyl- oder D-Tryptophylrest ist, und/oder der Aminosäurerest, der durch X<sub>2</sub> repräsentiert werden kann, ein Tyrosylrest ist, und/oder das Halogen, durch das der Rest des D-Aminosäurederivates, welches durch X<sub>1</sub> repräsentiert werden kann, und/oder der Rest des Aminosäurederivates, welcher durch X<sub>2</sub> repräsentiert werden kann, ringsubstituiert werden kann, Iod ist, wobei besonders bevorzugt der Rest des D-Aminosäurederivates, welcher durch X<sub>1</sub> repräsentiert werden kann, ein D-Diiodtyrosylrest ist, und/oder der Rest des Aminosäurederivates, welcher durch X<sub>2</sub> repräsentiert werden kann, ein Diiodtyrosylrest ist.  
 45
5. Verfahren gemäß den Ansprüchen 1 bis 3, **dadurch gekennzeichnet**, daß Octapeptid- oder Heptapeptidamidderivate hergestellt werden, in denen der Rest des Amins, welches durch A<sub>1</sub> repräsentiert werden kann, ein Indoliny- oder Phenylaminorest ist, die Alkoxygruppe, welche durch A<sub>1</sub> repräsentiert werden kann, eine solche mit 1 bis 4 Kohlenstoffatomen ist, die Aryloxygruppe, welche durch A<sub>1</sub> repräsentiert werden kann, eine Phenylloxygruppe ist, oder die Aralkyloxygruppe, welche durch A<sub>1</sub> repräsentiert werden kann, eine solche mit 1 bis 4 Kohlenstoffatomen in dem Alkylteil und einem Phenylteil als Arylteil ist.  
 50  
 55
6. Verfahren gemäß den Ansprüchen 1 bis 3, **dadurch gekennzeichnet**, daß Octapeptid- oder Heptapeptidamidderivate hergestellt werden, in denen die Alkylseitenkette des Aminosäurerestes oder seines Derivates, welcher durch X<sub>4</sub> repräsentiert werden kann, ein solcher mit 1 bis 4 Kohlenstoffatomen ist,

oder die Alkylseitenkette des Aminosäurerestes oder seines Derivates, welcher durch  $X_4$  repräsentiert werden kann, eine solche mit 1 bis 4 Kohlenstoffatomen im Alkylteil und einem Phenylteil als Arylteil ist, wobei insbesondere der Aminosäurerest, welcher durch  $X_4$  repräsentiert werden kann, ein Leucyl-, Valyl-, Isoleucyl-, Norvalyl-, Norleucyl-, Alanyl- oder Serylrest ist, und/oder der aromatische Aminosäurerest, welcher durch  $X_5$  repräsentiert werden kann, ein Tryptophyl- oder Phenylalanylrest ist, oder das Halogen, durch das der Rest des Derivats der aromatischen Aminosäure, welche durch  $X_5$  repräsentiert werden kann, substituiert werden kann, Iod ist, wobei besonders bevorzugt der Rest des Derivats der aromatischen Aminosäure, welche durch  $X_5$  repräsentiert werden kann, ein Tyrosyl- oder Diiodtyrosylrest ist, und/oder die  $C_{1-4}$ -Alkylgruppe, welche durch  $A_2$  repräsentiert werden kann, eine solche mit 1 oder 2 Kohlenstoffatomen ist, oder das Halogen, durch das die  $C_{1-4}$ -Alkylgruppe, welche durch  $A_2$  repräsentiert werden kann, substituiert werden kann, Fluor ist und/oder n 3 oder 4 ist.

7. Verfahren gemäß den Ansprüchen 1 bis 3, **dadurch gekennzeichnet**, daß Octapeptid- oder Heptapeptidamidderivate hergestellt werden, welche gewählt werden aus der aus



D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>,  
Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>,  
Pop-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub> und Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub> bestehenden Gruppe.


8. Verfahren zur Herstellung von Medikamenten, **dadurch gekennzeichnet**, daß als aktive Bestandteile 1 oder mehrere Verbindungen gemäß den Ansprüchen 1 bis 7 mit 1 oder mehreren Trägern und/oder Additiven, wie sie üblicherweise in der pharmazeutischen Technik verwendet werden, in einer per se bekannten Weise vermischt werden.



## Revendications


Revendications pour les Etats contractants suivants : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, PT, SE

- 5 1. Dérivés octapeptidyl ou heptapeptidyl amide de formule générale :




dans laquelle :

- $X_1$  représente un reste D-aminoacide aromatique ou un dérivé de celui-ci substitué sur le cycle par un halogène et/ou un groupe hydroxyle; ou un groupe D-phénylglycyle, p-hydroxyphénylglycyle, o-aminobenzoyl, m-aminobenzoyl, D-tétrahydroisoquinoléylcarbonyl, ou sarcosyl-analyte; ou un groupe :




dans lequel :

- $A_1$  représente le reste d'une amine aromatique, hétérocyclique, ou aliphatique comportant un groupe amine primaire ou secondaire, ou
- représente un groupe amine aromatique, hétérocyclique, ou aliphatique, primaire ou secondaire,
- ou
- représente un groupe alcoxy, aryloxy, ou aralkyloxy;
- $X_2$  représente un reste aminoacide aromatique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène et/ou un groupe hydroxyle; ou un groupe histidyle;
- $X_3$  représente un groupe D-tryptophyle, o-aminobenzoyl, m-aminobenzoyl, aspartyle, D-aspartyle,  $\beta$ -aspartyle,  $\beta$ -D-aspartyle, ou aminosuccinyle; ou un groupe :



dans lequel :

- $A_1$  est tel que défini ci-dessus;
- $X_4$  représente un reste aminoacide portant une chaîne latérale alkyle ou aralkyle, ou un dérivé de celui-ci, substitué par un groupe hydroxyle ou méthyle en position  $\beta$ ; ou un groupe prolyle ou  $\beta$ -alanyle; ou une liaison de valence;
- $X_5$  représente un reste aminoacide hétérocyclique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène ou un groupe hydroxyle; ou un groupe thréonyl;
- $R_1$  représente un atome d'hydrogène ou un groupe :



dans lequel :

- $A_2$  représente un atome d'hydrogène ou un groupe alkyle  $C_{1-4}$ , éventuellement substitué par un halogène;
- $R_2$  représente un atome d'hydrogène, un groupe -S-, ou un groupe -S-acétamidométhyle, phényle, ou p-hydroxyphényle;
- $R_3$  représente un atome d'hydrogène ou un groupe hydroxyle;

- R<sub>4</sub> représente un groupe phényle ou p-hydroxyphényle; ou un groupe -S-, ou -S-acétamidométhyle lorsque R<sub>2</sub> représente le même groupe; ou un groupe méthyle lorsque R<sub>3</sub> est un groupe hydroxyle, dans le cas où R<sub>2</sub> et R<sub>4</sub> sont chacun un groupe -S-, et une liaison de l'un est unie à une liaison de l'autre en une liaison unique représentant un pont entre R<sub>2</sub> et R<sub>4</sub>, et
- 5 - n est égal à 1, 2, 3, ou 4
- à condition que :
- a) si
- X<sub>1</sub> représente un groupe D-β-Nal ou D-Phe, éventuellement substitué par un halogène ou un groupe hydroxy,
- 10 X<sub>2</sub> représente un groupe Tyr ou Phe
- X<sub>3</sub> représente un groupe D-Trp
- X<sub>5</sub> représente un groupe Trp ou Thr, et
- R<sub>2</sub> et R<sub>4</sub> représentent chacun un groupe -S-, une liaison de l'un étant unie à une liaison de l'autre en une liaison unique représentant un pont entre R<sub>2</sub> et R<sub>4</sub>
- 15 X<sub>4</sub> est un groupe différent de Val, Thr, ou Ser,
- b) si
- X<sub>4</sub> représente un groupe Val, ou Thr, ou Ser, et
- X<sub>1</sub> représente un résidu D-aminoacide aromatique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène ou un groupe hydroxy,
- 20 X<sub>2</sub> représente un reste aminoacide aromatique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène ou un groupe hydroxy,
- au moins une des conditions suivantes soit remplie :
- α) X<sub>3</sub> est un groupe différent de D-Trp, ou
- β) R<sub>2</sub> et R<sub>4</sub> ensemble, représentent un groupe différent d'un groupe -S-S- (pont disulfure),
- 25 et les sels d'addition d'acide de ceux-ci.
2. Dérivés octapeptidyl ou heptapeptidyl amide selon la revendication 1, caractérisés en ce que le reste D-aminoacide pouvant être représenté par X<sub>1</sub>, est un reste D-phénylalanyle, D-naphtylalanyle, ou D-tryptophyle, et/ou le reste aminoacide pouvant être représenté par X<sub>2</sub> est un reste tyrosyle, et/ou
- 30 l'halogène par lequel le reste du dérivé aminoacide pouvant être représenté par X<sub>1</sub>, et/ou le reste du dérivé aminoacide pouvant être représenté par X<sub>2</sub>, peuvent être substitués sur le cycle, est l'iode, par préférence particulière, le reste de dérivé D-aminoacide pouvant être représenté par X<sub>1</sub>, étant un reste D-diiodotyrosyle, et/ou le reste, de dérivé aminoacide pouvant être représenté par X<sub>2</sub>, étant un
- 35 reste diiodotyrosyle.
3. Dérivés octapeptidyl ou heptapeptidyl amide selon la revendication 1, caractérisés en ce que le reste de l'amine pouvant être représenté par A<sub>1</sub>, est un reste indolyle ou phénylamino, le groupe alcoxy pouvant être représenté par A<sub>1</sub>, est tel qu'il comporte 1 à 4 atome(s) de carbone, le groupe aryloxy pouvant être représenté par A<sub>1</sub>, est un groupe phényloxy, ou le groupe aralkyloxy pouvant être
- 40 représenté par A<sub>1</sub>, est tel qu'il comporte 1 à 4 atome(s) de carbone dans la partie alkyle, et qu'il a une partie phényle en tant que partie aryle.
4. Dérivés octapeptidyl ou heptapeptidyl amide selon les revendications 1 à 3, caractérisés en ce que la chaîne latérale alkyle du reste aminoacide ou de son dérivé, pouvant être représentée par X<sub>4</sub> est telle
- 45 qu'elle comporte 1 à 4 atome(s) de carbone, ou que la chaîne latérale aralkyle du reste aminoacide ou de son dérivé, pouvant être représentée par X<sub>4</sub>, est telle qu'elle comporte 1 à 4 atome(s) de carbone dans la partie alkyle, et qu'elle a une partie phényle en tant que partie aryle, et en particulier, le reste aminoacide pouvant être représenté par X<sub>4</sub> est un reste leucyle, valyle, isoleucyle, norvalyle, norleucyle
- 50 alanyle ou séryle, et/ou le reste aminoacide pouvant être représenté par X<sub>5</sub> est un reste tryptophyle ou phénylalanyle, ou l'halogène par lequel le reste du dérivé aminoacide aromatique pouvant être représenté par X<sub>5</sub>, peut être substitué, est l'iode, et, par préférence particulière, le reste de dérivé aminoacide aromatique pouvant être représenté par X<sub>5</sub>, est un reste tyrosyle ou diiodotyrosyle, et/ou le
- groupe alkyle C<sub>1-4</sub> pouvant être représenté par A<sub>2</sub>, est tel qu'il comporte 1 à 2 atomes de carbone, ou
- 55 l'halogène par lequel le groupe alkyle C<sub>1-4</sub>, pouvant être représenté par A<sub>2</sub>, peut être substitué, est le fluor, et/ou n est égal à 3 ou à 4.

5. Composé selon les revendications 1 à 4, choisi au sein du groupe comprenant les groupes :

-  $\beta$ -aspartyl-(indolyl)-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub>

- D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr-NH<sub>2</sub>

- D-Phe-Cys-Tyr-D-Trp-Lys-Leu-Cys-Thr-NH<sub>2</sub>

- D-Phe-Cys-Tyr-β-Asp(indolinyI)-Lys-Leu-Cys-Thr-NH<sub>2</sub>

- D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>

- Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>

- Pop-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>, et

- Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub>.

6. Procédé pour la préparation des composés selon les revendications 1 à 5, caractérisé en ce que, lors de l'utilisation de la méthode de synthèse des polypeptides en phase solide, les aminoacides protégés sont graduellement condensés sur le véhicule résine solide par couplage au carbodiimide ou ester actif, dans l'ordre correspondant, puis le produit final est séparé du véhicule solide par clivage acide ou basique, et les groupes protecteurs sont séparés des aminoacides, avant, pendant, ou après le clivage à partir de la résine, et, si désiré, le dérivé octapeptidyl ou heptapeptidyl amide de formule générale I ainsi obtenu est converti en un sel d'addition d'acide, en le faisant réagir avec un acide et/ou si désiré, la base libre est libérée à partir du sel d'addition d'acide, en le faisant réagir avec une base, et, si désiré, cette dernière est convertie en un autre sel d'addition d'acide, en la faisant réagir avec un acide.

7. Procédé selon à revendication 6, caractérisé en ce qu'une résine benzhydrylamine, ou une résine polystyrène chlorométhylée est utilisée comme véhicule résine solide.


8. Procédé selon les revendications 6 ou 7, caractérisé en ce que le produit final contenant les groupes protecteurs est scindé par le gaz fluorhydrique et/ou par ammonolyse.

9. Médicaments caractérisés en ce qu'ils contiennent comme ingrédient(s) actif(s), un ou plusieurs composés selon les revendications 1 à 8, convenablement, en mélange avec un ou plusieurs véhicule(s), et/ou des additifs communément utilisés en technique pharmaceutique.

10. Utilisation des composés selon les revendications 1 à 5, pour la préparation de médicaments, en particulier pour le traitement des mammifères, y compris l'homme, souffrant de maladie tumorale, et/ou pour l'inhibition de leur activité tyrosine kinase.

**Revendications pour les Etats c ntractants suivants : ES, GR**

1. Procédé pour la préparation de dérivés octapeptidyl ou heptapeptidyl amide de formule générale :




55 dans laqu lle :

- X<sub>1</sub> représente un reste D-aminoacide aromatique, ou un dérivé de celui-ci substitué sur le cycle par un halogène et/ou un groupe hydroxyle; ou un groupe D-phénylglycyle, p-hydroxyphénylglycyle, o-aminobenzoyl, m-aminobenzoyl, D-tétrahydroisoquinoléylcarbonyl, ou sarcosylanalyle;

ou un groupe :

5




10

dans lequel :

- 15  $\text{A}_1$  représente le reste d'une amine aromatique, hétérocyclique, ou aliphatique comportant un groupe amine primaire ou secondaire, ou
- représente un groupe amine aromatique, hétérocyclique, ou aliphatique, primaire ou secondaire,
- ou
- représente un groupe alcoxy, aryloxy, ou aralkyloxy;
- 20 -  $\text{X}_2$  représente un reste aminoacide aromatique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène et/ou un groupe hydroxyle; ou un groupe histidyle;
- $\text{X}_3$  représente un groupe D-tryptophyle, o-aminobenzoyl, m-aminobenzoyl, aspartyle, D-aspartyle,  $\beta$ -aspartyle,  $\beta$ -D-aspartyle, ou aminosuccinyle; ou un groupe :

25




30

dans lequel :

- 35  $\text{A}_1$  est tel que défini ci-dessus;
- $\text{X}_4$  représente un reste aminoacide portant une chaîne latérale alkyle ou aralkyle, ou un dérivé de celui-ci, substitué par un groupe hydroxyle ou méthyle en position  $\beta$ ; ou un groupe prolyle ou  $\beta$ -alanyle; ou une liaison de valence;
- 40 -  $\text{X}_5$  représente un reste aminoacide hétérocyclique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène ou un groupe hydroxyle; ou un groupe thréonyl;
- $\text{R}_1$  représente un atome d'hydrogène ou un groupe :

45



50

dans lequel :

- $\text{A}_2$  représente un atome d'hydrogène ou un groupe alkyle  $\text{C}_{1-4}$ , éventuellement substitué par un halogène;
- $\text{R}_2$  représente un atome d'hydrogène, un groupe -S-, ou un groupe -S-acétamidométhyle, phényle, ou p-hydroxyphényle;
- 55 -  $\text{R}_3$  représente un atome d'hydrogène ou un groupe hydroxyle;
- $\text{R}_4$  représente un groupe phényle ou p-hydroxyphényle; ou un groupe -S-, ou -S-acétamidométhyle lorsque  $\text{R}_2$  représente le même groupe; ou un groupe méthyle lorsque  $\text{R}_3$  est un groupe hydroxyle, dans le cas où  $\text{R}_2$  et  $\text{R}_4$  sont chacun un groupe -S-, et une liaison de l'un est unie à

une liaison de l'autre en une liaison unique représentant un pont entre R<sub>2</sub> et R<sub>4</sub>, et  
 - n est égal à 1, 2, 3 ou 4,  
 à condition que :

a) si

- 5 X<sub>1</sub> représente un groupe D-β-Nal ou D-Phe, éventuellement substitué par un halogène ou un groupe hydroxy,  
 X<sub>2</sub> représente un groupe Tyr ou Phe  
 X<sub>3</sub> représente un groupe D-Trp  
 X<sub>5</sub> représente un groupe Trp ou Thr, et  
 10 R<sub>2</sub> et R<sub>4</sub> représentent chacun un groupe -S-, une liaison de l'un étant unie à une liaison de l'autre en une liaison unique représentant un pont entre R<sub>2</sub> et R<sub>4</sub> est un groupe différent de Val, Thr, ou Ser,

b) si

- 15 X<sub>4</sub> représente un groupe Val, ou Thr, ou Ser, et  
 X<sub>1</sub> représente un résidu D-aminoacide aromatique, ou un dérivé de celui-ci, substitué sur le cycle par un halogène ou un groupe hydroxy,  
 X<sub>2</sub> représente un reste aminoacide aromatique, ou un dérivé de celui-ci substitué sur le cycle par un halogène ou un groupe hydroxy,

au moins une des conditions suivantes soit remplie :

- 20 α) X<sub>3</sub> est un groupe différent de D-Trp, ou  
 β) R<sub>2</sub> et R<sub>4</sub> ensemble, représentent un groupe différent d'un groupe -S-S- (pont disulfure), et les sels d'addition d'acide de ceux-ci,  
 caractérisé en ce que, lors de l'utilisation de la méthode de synthèse des polypeptides en phase  
 25 solide, les aminoacides protégés sont graduellement condensés sur le véhicule résine solide par couplage au carbodiimide ou ester actif dans l'ordre correspondant, puis le produit final est séparé du véhicule solide par clivage acide ou basique, et les groupes protecteurs sont séparés des aminoacides, avant, pendant, ou après le clivage à partir de la résine, et, si désiré, le dérivé octapeptidyl ou heptapeptidyl amide de formule générale I ainsi obtenu est converti en un sel d'addition d'acide, en le  
 30 faisant réagir avec un acide et/ou si désiré, la base libre est libérée à partir du sel d'addition d'acide, en le faisant réagir avec une base, et, si désiré, cette dernière est convertie en un autre sel d'addition d'acide, en la faisant réagir avec un acide.

2. Procédé selon la revendication 1, caractérisé en ce qu'une résine benzhydramine, ou une résine polystyrène chlorométhylée est utilisée comme véhicule résine solide.

3. Procédé selon les revendications 1 ou 2, caractérisé en ce que le produit final contenant les groupes protecteurs est scindé par le gaz fluorhydrique et/ou par ammonolyse.





4. Procédé selon les revendications 1 à 3, pour la préparation de dérivés octapeptidyl ou heptapeptidyl amide, caractérisé en ce que le reste D-aminoacide pouvant être représenté par X<sub>1</sub>, est un reste D-phénylalanyle, D-naphtylalanyle, ou D-tryptophyle, et/ou le reste aminoacide pouvant être représenté par X<sub>2</sub> est un reste tyrosyle, et/ou l'halogène par lequel le reste du dérivé D-aminoacide pouvant être représenté par X<sub>1</sub>, et/ou le reste du dérivé aminoacide pouvant être représenté par X<sub>2</sub>, peuvent être substitués sur le cycle, est l'iode, par préférence particulière, le reste du dérivé D-aminoacide pouvant être représenté par X<sub>1</sub>, étant un reste D-diiodotyrosyle, et/ou le reste de dérivé aminoacide pouvant être représenté par X<sub>2</sub>, étant un reste diiodotyrosyle.

5. Procédé selon les revendications 1 à 3, pour la préparation de dérivés octapeptidyl ou heptapeptidyl amide, caractérisé en ce que le reste de l'amine pouvant être représenté par A<sub>1</sub>, est un reste indolinyloxy ou phénylaminio, le groupe alcoxy pouvant être représenté par A<sub>1</sub>, est tel qu'il comporte 1 à 4 atome(s) de carbone, le groupe aryloxy pouvant être représenté par A<sub>1</sub>, est un groupe phényloxy, ou le groupe aralkyloxy pouvant être représenté par A<sub>1</sub>, est tel qu'il comporte 1 à 4 atome(s) de carbone dans la partie alkyle, et qu'il a une partie phényle en tant que partie aryle.

6. Procédé selon les revendications 1 à 3, pour la préparation de dérivés octapeptidyl ou heptapeptidyl amide, caractérisé en ce que la chaîne latérale alkyle du reste aminoacide ou de son dérivé, pouvant être représentée par X<sub>4</sub>, est telle qu'elle comporte 1 à 4 atome(s) de carbone, ou que la chaîne latérale aralkyle du reste aminoacide ou de son dérivé, pouvant être représentée par X<sub>4</sub>, est telle qu'elle

comporte 1 à 4 atome(s) de carbone dans la partie alkyle, et qu'elle a une partie phényle en tant que partie aryle, et en particulier, le reste aminoacide pouvant être représenté par  $X_4$  est un reste leucyle, valyle, isoleucyle, norvalyle, norleucyle, alanyle ou séryle, et/ou le reste aminoacide pouvant être représenté par  $X_5$  est un reste tryptophyle ou phénylalanyle, ou l'halogène par lequel le reste du dérivé aminoacide aromatique pouvant être représenté par  $X_5$ , peut être substitué, est l'iode, et, par préférence particulière, le reste de dérivé aminoacide aromatique pouvant être représenté par  $X_5$ , est un reste tyrosyle ou diiodotyrosyle, et/ou le groupe alkyle  $C_{1-4}$  pouvant être représenté par  $A_2$ , est tel qu'il comporte 1 à 2 atomes de carbone, ou l'halogène par lequel le groupe alkyle  $C_{1-4}$ , pouvant être représenté par  $A_2$ , peut être substitué, est le fluor et/ou n est égal à 3 ou à 4.

7. Procédé selon les revendications 1 à 3, pour la préparation de dérivés octapeptidyl ou heptapeptidyl amide, choisis au sein du groupe comprenant les groupes :



- D-Phe-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>  
 - Ac-Sar-Ala-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>  
 - Pop-Tyr-Tyr-Aa-Lys-Val-Phe-Trp-NH<sub>2</sub>, et  
 - Aa-Cys(Acm)-Tyr-D-Trp-Lys-Val-Cys(Acm)-Trp-NH<sub>2</sub>.

8. Procédé de préparation de médicaments caractérisé par le mélange, en tant qu'ingrédient(s) actif(s), d'un ou de plusieurs composés préparés par le procédé selon les revendications 1 à 7, avec un ou plusieurs véhicule(s) et/ou additif(s) communément utilisés en technique pharmaceutique d'une manière connue per se.